antibodies -online.com





UGT1A10 Protein (Myc-DYKDDDDK Tag)






Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Overview		
Quantity:	20 μg	
Target:	UGT1A10	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This UGT1A10 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	 Recombinant human UGT1A10 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	UGT1A10	
Alternative Name:	Ugt1a10 (UGT1A10 Products)	
Background:	This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes.	
	several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate	

Target Details

	Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. The enzyme encoded by this gene has glucuronidase activity on mycophenolic acid, coumarins, and quinolines.		
Molecular Weight:	57.2 kDa		
NCBI Accession:	NP_061948		
Pathways:	Steroid Hormone Biosynthesis, Regulation of Lipid Metabolism by PPARalpha		
Application Details			
Application Notes:	Recombinant human proteins can be used for:		
	Native antigens for optimized antibody production		
	Positive controls in ELISA and other antibody assays		
Comment:	The tag is located at the C-terminal.		
Restrictions:	For Research Use only		
Handling			
Concentration:	50 μg/mL		
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.		
Storage:	-80 °C		
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.		



Western Blotting

Image 1. Validation with Western Blot