## antibodies -online.com





## **UGT1A4 Protein (Myc-DYKDDDDK Tag)**



Image




Go to Product page

( )	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	1 V I	ew

Overview		
Quantity:	20 μg	
Target:	UGT1A4	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This UGT1A4 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	<ul> <li>Recombinant human UGT1A4 protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	UGT1A4	
Alternative Name:	Ugt1a4 (UGT1A4 Products)	
Background:	This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway	
	that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into	
	water-soluble, excretable metabolites. This gene is part of a complex locus that encodes	
	several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons	
	followed by four common exons. Four of the alternate first exons are considered pseudogenes.	

## **Target Details**

	Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. This enzyme has some glucuronidase activity towards bilirubin, although is is more active on amines, steroids, and sapogenins.	
Molecular Weight:	57 kDa	
NCBI Accession:	NP_009051	
Pathways:	Steroid Hormone Biosynthesis, Regulation of Lipid Metabolism by PPARalpha	
Application Details		
Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	
Handling		
Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
torage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freez immediately. Only 2-3 freeze thaw cycles are recommended.		



## **Western Blotting**

Image 1. Validation with Western Blot