antibodies .- online.com





VRK3 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)





Go to Product page

| \sim | | |
|---------|--------|----------------|
| 0ve | ı۱/۱۲د | $\triangle VV$ |
| \circ | ا ۷ ار | CVV |

| Overview | |
|-------------------------------|--|
| Quantity: | 20 μg |
| Target: | VRK3 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This VRK3 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human VRK3 (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | VRK3 |
| Alternative Name: | Vrk3 (VRK3 Products) |
| Background: | This gene encodes a member of the vaccinia-related kinase (VRK) family of serine/threonine |
| | protein kinases. In both human and mouse, this gene has substitutions at several residues |
| | within the ATP binding motifs that in other kinases have been shown to be required for |
| | catalysis. In vitro assays indicate the protein lacks phosphorylation activity. The protein, |
| Alternative Name: | Vrk3 (VRK3 Products) This gene encodes a member of the vaccinia-related kinase (VRK) family of serine/threonin protein kinases. In both human and mouse, this gene has substitutions at several residues within the ATP binding motifs that in other kinases have been shown to be required for |

Target Details

| | however, likely retains its substrate binding capability. This gene is widely expressed in human |
|-------------------|--|
| | tissues and its protein localizes to the nucleus. Alternative splicing results in multiple |
| | transcripts encoding different isoforms. |
| Molecular Weight: | 52.7 kDa |
| NCBI Accession: | NP_057524 |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot