# antibodies - online.com





## WASP Protein (Myc-DYKDDDDK Tag)



Image



$\sim$				
	$ V \cap$	r\/I	19	٨

Overview	
Quantity:	20 μg
Target:	WASP (WAS)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This WASP protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human WAS / IMD2 protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	WASP (WAS)
Alternative Name:	Was,imd2 (WAS Products)
Background:	The Wiskott-Aldrich syndrome (WAS) family of proteins share similar domain structure, and are
	involved in transduction of signals from receptors on the cell surface to the actin cytoskeleton.
	The presence of a number of different motifs suggests that they are regulated by a number of
	different stimuli, and interact with multiple proteins. Recent studies have demonstrated that
	these proteins, directly or indirectly, associate with the small GTPase, Cdc42, known to regulate

#### **Target Details**

formation of actin filaments, and the cytoskeletal organizing complex, Arp2/3. Wiskott-Aldrich syndrome is a rare, inherited, X-linked, recessive disease characterized by immune dysregulation and microthrombocytopenia, and is caused by mutations in the WAS gene. The WAS gene product is a cytoplasmic protein, expressed exclusively in hematopoietic cells, which show signalling and cytoskeletal abnormalities in WAS patients. A transcript variant arising as a result of alternative promoter usage, and containing a different 5' UTR sequence, has been described, however, its full-length nature is not known.

Molecular Weight:	52.7 kDa
-------------------	----------


NCBI Accession: NP\_000368

#### **Application Details**

Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	

#### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



### **Western Blotting**

Image 1. Validation with Western Blot