

## Datasheet for ABIN2735603

## **WISP3 Protein (Transcript Variant 1)**



Go to Product page

| _ |       |             |    |    |             |     |
|---|-------|-------------|----|----|-------------|-----|
|   | V     | $\triangle$ | r۱ | /1 | $\triangle$ | Λ/  |
|   | ' V ' |             | ΙV |    |             | v v |

| Quantity:                    | 20 μg  |
|------------------------------|--|
| Target:                      | WISP3  |
| Protein Characteristics:     | Transcript Variant 1   |
| Origin:                      | Human  |
| Source:                      | Escherichia coli (E. coli)   |
| Protein Type:                | Recombinant  |
| Biological Activity:         | Active   |
| Application:                 | Functional Studies (Func), Antibody Production (AbP), Protein Interaction (PI), Standard (STD)   |
| Product Details              |  |
| Specificity:                 | Optimal preservation of protein structure, post-translational modifications and functions.   |
| Characteristics:             | <ul> <li>Recombinant human WISP3 (transcript variant 1) protein expressed in E. coli.</li> <li>Produced with end-sequenced ORF clone</li> <li>Tested for bioactivity.</li> </ul> |
| Purity:                      | > 95 % as determined by SDS-PAGE and Coomassie blue staining   |
| Endotoxin Level:             | Endotoxin level is <0.1 ng/μg of protein (<1EU/μg).  |
| Biological Activity Comment: | ED50 was determined by the dose-dependant proliferation of the MCF-7 cell line. The expected ED50 for this effect is 0.2-0.3 ug/ml.  |
| Target Details               |  |
| Target:                      | WISP3  |
|                              |  |

## **Target Details**

| Alternative Name:   | Wisp3 (WISP3 Products)  |  |  |  |  |
|---------------------|---|--|--|--|--|
| Background:         | This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein   |  |  |  |  |
|                     | subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a  |  |  |  |  |
|                     | member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse   |  |  |  |  |
|                     | developmental processes. The CTGF family members are characterized by four conserved  |  |  |  |  |
|                     | cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C  |  |  |  |  |
|                     | module, thrombospondin domain and C-terminal cystine knot-like domain. This gene is   |  |  |  |  |
|                     | overexpressed in colon tumors. It may be downstream in the WNT1 signaling pathway that is   |  |  |  |  |
|                     | relevant to malignant transformation. Mutations of this gene are associated with progressive  |  |  |  |  |
|                     | pseudorheumatoid dysplasia, an autosomal recessive skeletal disorder, indicating that the gene  |  |  |  |  |
|                     | is essential for normal postnatal skeletal growth and cartilage homeostasis. Multiple transcript  |  |  |  |  |
|                     | variants encoding different isoforms have been found for this gene.   |  |  |  |  |
| Molecular Weight:   | 36.8 kDa  |  |  |  |  |
| NCBI Accession:     | NP_003871   |  |  |  |  |
| Pathways:           | WNT Signaling, Growth Factor Binding  |  |  |  |  |
| Application Details |   |  |  |  |  |
| Application Notes:  | Recombinant human proteins can be used for:   |  |  |  |  |
|                     | Native antigens for optimized antibody production   |  |  |  |  |
|                     | Positive controls in ELISA and other antibody assays  |  |  |  |  |
|                     | Protein-protein interaction   |  |  |  |  |
|                     | In vitro biochemical assays and cell-based functional assays  |  |  |  |  |
| Restrictions:       | For Research Use only   |  |  |  |  |
| Handling            |   |  |  |  |  |
| Buffer:             | Lyophilized from a 0.2 µM filtered solution of 20 mM phosphate buffer,100 mM NaCl, pH 7.2   |  |  |  |  |
| Handling Advice:    | Resuspend the protein in the desired concentration in proper buffer   |  |  |  |  |
| Storage:            | -80 °C  |  |  |  |  |
| Storage Comment:    | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |  |  |  |  |