# antibodies -online.com





## WNK4 Protein (Myc-DYKDDDDK Tag)



Image



| $C \circ + \circ$ | Drad | 110+ | 2000 |
|-------------------|------|------|------|
| Go to             | PIOU | uCt  | Daue |

| Overview                      |  |
|-------------------------------|--|
| Quantity:                     | 20 μg  |
| Target:                       | WNK4   |
| Origin:                       | Human  |
| Source:                       | HEK-293 Cells  |
| Protein Type:                 | Recombinant  |
| Purification tag / Conjugate: | This WNK4 protein is labelled with Myc-DYKDDDDK Tag.   |
| Application:                  | Antibody Production (AbP), Standard (STD)  |
| Product Details               |  |
| Characteristics:              | <ul> <li>Recombinant human WNK4 protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>   |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |
| Target Details                |  |
| Target:                       | WNK4   |
| Alternative Name:             | Wnk4 (WNK4 Products)   |
| Background:                   | This gene encodes a member of the WNK family of serine-threonine protein kinases. The kinase is part of the tight junction complex in kidney cells, and regulates the balance between NaCl reabsorption and K(+) secretion. The kinase regulates the activities of several types of ion channels, cotransporters, and exchangers involved in electrolyte flux in epithelial cells. |

Mutations in this gene result in pseudohypoaldosteronism type IIB.[provided by RefSeq, Sep

### Target Details

|                   | 2009].    |
|-------------------|-----------|
| Molecular Weight: | 134.6 kDa |
| NCBI Accession:   | NP_115763 |


## **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |
|--------------------|--|
|                    | Native antigens for optimized antibody production    |
|                    | Positive controls in ELISA and other antibody assays |
| Comment:           | The tag is located at the C-terminal.                |
| Restrictions:      | For Research Use only                                |

### Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

#### **Images**



#### **Western Blotting**

Image 1. Validation with Western Blot