Datasheet for ABIN2735608

# WNT16 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)

**1** Image

Go to Product page

## Overview

| | |
|---|---|
| Quantity: | 20 µg |
| Target: | WNT16 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This WNT16 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |

## Product Details

| | |
|---|---|
| Characteristics: | • Recombinant human WNT16 (transcript variant 1) protein expressed in HEK293 cells.<br>• Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |

## Target Details

| | |
|---|---|
| Target: | WNT16 |
| Alternative Name: | Wnt16 (WNT16 Products) |
| Background: | The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It contains two transcript variants diverging at the 5' termini. |

## Target Details

These two variants are proposed to be the products of separate promoters and not to be splice variants from a single promoter. They are differentially expressed in normal tissues, one of which (variant 2) is expressed at significant levels only in the pancreas, whereas another one (variant 1) is expressed more ubiquitously with highest levels in adult kidney, placenta, brain, heart, and spleen.

| | |
|---|---|
| Molecular Weight: | 37.6 kDa |
| NCBI Accession: | NP_476509 |
| Pathways: | WNT Signaling |

## Application Details

| | |
|---|---|
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

## Handling

| | |
|---|---|
| Concentration: | 50 µg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

# Images



## Western Blotting

**Image 1.** Validation with Western Blot