

Datasheet for ABIN2745570

anti-TEK antibody (Extracellular Domain)[Go to Product page](#)**1** Image

Overview

| | |
|----------------------|---|
| Quantity: | 100 µg |
| Target: | TEK |
| Binding Specificity: | Extracellular Domain |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Application: | Western Blotting (WB), ELISA, Flow Cytometry (FACS) |

Product Details

| | |
|-------------------|---|
| Immunogen: | Recombinant human soluble extracellular domain of TIE-2. |
| Clone: | Tek2 |
| Isotype: | IgG1 |
| Specificity: | Native human TIE-2 in ELISA experiments and on the surface of different human cell types. |
| Cross-Reactivity: | Human |
| Purification: | Protein G purified. |

Target Details

| | |
|-------------------|---|
| Target: | TEK |
| Alternative Name: | TIE-2 (TEK Products) |
| Background: | TIE-2 acts as cell-surface receptor for ANGPT1, ANGPT2 and ANGPT4 and regulates |

Target Details

angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, reorganization of the actin cytoskeleton, but also maintenance of vascular quiescence. It has anti-inflammatory effects by preventing the leakage of proinflammatory plasma proteins and leukocytes from blood vessels. It is required for normal angiogenesis and heart development during embryogenesis and for post-natal hematopoiesis. After birth, it activates or inhibits angiogenesis, depending on the context. It inhibits angiogenesis and promotes vascular stability in quiescent vessels, where endothelial cells have tight contacts. In migrating endothelial cells that lack cell-cell adhesions, ANG1 recruits TEK to contacts with the extracellular matrix, leading to the formation of focal adhesion complexes, activation of PTK2/FAK and of the downstream kinases MAPK1/ERK2 and MAPK3/ERK1 and ultimately to the stimulation of sprouting angiogenesis.


| | |
|-----------|--|
| UniProt: | Q02763 |
| Pathways: | RTK Signaling, Growth Factor Binding |

Application Details

| | |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|--|
| Format: | Lyophilized |
| Reconstitution: | Centrifuge vial prior to opening. Reconstitute with sterile water to a concentration of 0.1-1.0 mg/mL. |
| Concentration: | Lot specific |
| Buffer: | Lyophilized. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Short Term Storage: +4°C Long Term Storage: -20°C Stable for at least 6 months after receipt when stored at -20°C. |
| Expiry Date: | 6 months |



Immunohistochemistry

Image 1. Immunohistochemistry detection of endogenous TIE-2 in cryo sections of human spleen using anti-TIE-2 (human), mAb (tek2) .