

[Go to Product page](#)

Datasheet for ABIN2747908

Src ELISA Kit

Overview

Quantity:	96 tests
Target:	Src
Binding Specificity:	pTyr419, total
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human Phospho-SRC (Y419) and Total SRC ELISA Kit. This assay semi-quantitatively measures phosphorylated SRC (Tyr419) and Total SRC in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Human SRC phosphorylated at site Tyrosin-419 and Total SRC
Characteristics:	<ul style="list-style-type: none">• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)• Screen numerous different cell lysates without performing a Western Blot analysis• Minimal hands-on time, convenient, and non-radioactive material
Components:	<ul style="list-style-type: none">• Pre-Coated 96-well Strip Microplate• Wash Buffer

Product Details

- Anti-Phospho Antibody
- Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- Streptavidin-Conjugated HRP
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	Src
Alternative Name:	SRC (Src Products)
Target Type:	Viral Protein
Background:	Phospho-SRC (Y419). Full Name = Proto-oncogene tyrosine-protein kinase Src (P12931)
Gene ID:	6714
UniProt:	P12931
Pathways:	JAK-STAT Signaling , Neurotrophin Signaling Pathway , Intracellular Steroid Hormone Receptor Signaling Pathway , Regulation of Intracellular Steroid Hormone Receptor Signaling , Cellular Response to Molecule of Bacterial Origin , Cell-Cell Junction Organization , Regulation of Carbohydrate Metabolic Process , Autophagy , CXCR4-mediated Signaling Events , Signaling Events mediated by VEGFR1 and VEGFR2 , Smooth Muscle Cell Migration , Negative Regulation of intrinsic apoptotic Signaling , Platelet-derived growth Factor Receptor Signaling , Thromboxane A2 Receptor Signaling , Signaling of Hepatocyte Growth Factor Receptor , VEGF Signaling

Application Details

Sample Volume: 100 µL

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 µL of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 µL of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

Assay Procedure:

Prepare all reagents and samples as instructed in the manual.

Add 100 µL of sample or positive control to each well.

Incubate 2.5 h at RT or O/N at 4 °C.

Add 100 µL of prepared primary antibody to each well.

Incubate 1 h at RT.

Add 100 µL of prepared 1X HRP-Streptavidin to each well.

Incubate 1 h at RT.

Add 100 µL of TMB One-Step Substrate Reagent to each well.

Incubate 30 min at RT.

Add 50 µL of Stop Solution to each well.

Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment:

Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months