# antibodies -online.com





#### Datasheet for ABIN2747952

## **Angiopoietin 1 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	Angiopoietin 1 (ANGPT1)
Reactivity:	Human, Rabbit
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details	
Purpose:	Rabbit Angiopoietin-1 ELISA Kit for Serum, Plasma, and Cell Culture Supernatants.
Sample Type:	Plasma, Cell Culture Supernatant, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair recognizes Rabbit Angiopoietin-1.
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> </ul>

#### **Product Details**

- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 µL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- · Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

#### **Target Details**

Target:	Angiopoietin 1 (ANGPT1)
Alternative Name:	Angiopoietin-1 (ANGPT1 Products)
Background:	Angiopoietin-1 (ANGPT1)
UniProt:	G1SK98
Pathways:	RTK Signaling, Glycosaminoglycan Metabolic Process

Application Details	
Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 µL of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Assay Procedure:	Prepare all reagents, samples and standards as instructed in the manual.Add 100 μL of
	standard or sample to each well.Incubate 2.5 h at RT or O/N at 4 °C.Add 100 µL of prepared
	biotin antibody to each well.Incubate 1 h at RT.Add 100 µL of prepared Streptavidin solution to

### **Application Details**

each well.Incubate 45 min at RT.Add 100 μL of TMB One-Step Substrate Reagent to each well.Incubate 30 min at RT.Add 50 μL of Stop Solution to each well.Read at 450 nm
immediately.
"""Tediately.
For Research Use only
-20 °C
The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated
freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is
recommended to store at -80°C.
recommended to store at 700 C.
•