

# Datasheet for ABIN2748012

## **CD27 ELISA Kit**



#### Overview

Quantity:	96 tests
Target:	CD27
Reactivity:	Rat
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details	
Purpose:	Custom Rat CD27 (TNFRSF7) ELISA Kit.
Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Rat CD27 (TNFRSF7)
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> </ul>

### **Product Details**

- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 µL to 1 µL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

## **Target Details**

Target:	CD27
Alternative Name:	CD27 (CD27 Products)
Background:	CD27 (TNFRSF7)
Gene ID:	500318
UniProt:	Q501W2
Pathways:	Production of Molecular Mediator of Immune Response, Cancer Immune Checkpoints

## **Application Details**

Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 μL of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 $\mu$ L of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Assay Procedure:	Prepare all reagents, samples and standards as instructed in the manual.Add 100 µL of

## **Application Details**

Expiry Date:

6 months

Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.
Storage Comment:	-20 °C  The entire kit may be stored at 20°C for up to 1 year from the data of chipment. Avoid repeated
Handling	
Restrictions:	For Research Use only
	standard or sample to each well.Incubate 2.5 h at RT or O/N at 4 °C.Add 100 $\mu$ L of prepared biotin antibody to each well.Incubate 1 h at RT.Add 100 $\mu$ L of prepared Streptavidin solution to each well.Incubate 45 min at RT.Add 100 $\mu$ L of TMB One-Step Substrate Reagent to each well.Incubate 30 min at RT.Add 50 $\mu$ L of Stop Solution to each well.Read at 450 nm immediately.