

# Datasheet for ABIN2748182

### **IGF2 ELISA Kit**



### Overview

Quantity:	96 tests
Target:	IGF2
Reactivity:	Rhesus Monkey
Method Type:	Sandwich ELISA
Detection Range:	0.2-50 ng/mL
Minimum Detection Limit:	0.2 ng/mL
Application:	ELISA

#### **Product Details**

Troduct Details	
Purpose:	Rhesus Monkey IGF-2 ELISA Kit for Cell Culture Supernatants, Plasma, and Serum samples.
Sample Type:	Cell Culture Supernatant, Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Rhesus Macaque Insulin-like Growth Factor 2 (IGF2)
Sensitivity:	24 pg/mL
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>

### **Product Details**

#### Components:

- · Pre-Coated 96-well Strip Microplate
- · Wash Buffer
- · Stop Solution
- Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 μL to 1 μL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

### **Target Details**

Target:	IGF2
Alternative Name:	IGF-2 (IGF2 Products)
Background:	Gene Names: IFNG Protein names: Interferon gamma (IFN-gamma)
UniProt:	G7NBL8
Pathways:	Hormone Activity, Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process, Regulation of Carbohydrate Metabolic Process, Activated T Cell Proliferation

### **Application Details**

Application Notes:	Recommended Dilution for serum and plasma samples2 fold
Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	<ol> <li>Prepare all reagents, samples and standards as instructed in the manual.</li> <li>Add 100 μL of standard or sample to each well.</li> <li>Incubate 2.5 h at RT or O/N at 4 °C.</li> </ol>

## **Application Details**

	4. Add 100 μL of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated
	freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is
	recommended to store at -80°C.
Expiry Date:	6 months