



[Go to Product page](#)

Datasheet for ABIN2748349
CCL4 ELISA Kit

Overview

Quantity: 96 tests

Target: CCL4

Reactivity: Rhesus Monkey

Method Type: Sandwich ELISA

Application: ELISA

Product Details

Purpose: Custom Rhesus Macaque MIP-1 beta (CCL4) ELISA Kit.

Sample Type: Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Lysate

Analytical Method: Quantitative

Detection Method: Colorimetric

Specificity: The antibody pair provided in this kit recognizes Rhesus Macaque MIP-1 beta (CCL4)

Characteristics:

- Strip plates and additional reagents allow for use in multiple experiments
- Quantitative protein detection
- Establishes normal range
- The best products for confirmation of antibody array data

Components:

- Pre-Coated 96-well Strip Microplate
- Wash Buffer
- Stop Solution
- Assay Diluent(s)
- Lyophilized Standard
- Biotinylated Detection Antibody

Product Details

- Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 μ L to 1 μ L volumes
- Adjustable 1-25 μ L pipettes for reagent preparation
- 100 μ L and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

Target Details

Target: CCL4

Alternative Name: MIP-1 beta ([CCL4 Products](#))

Background: MIP-1 beta (CCL4)

Gene ID: 574217

UniProt: [Q8HYQ2](#)

Application Details

Sample Volume: 100 μ L

Plate: Pre-coated

Protocol:

1. Prepare all reagents, samples and standards as instructed in the manual.
2. Add 100 μ L of standard or sample to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 μ L of prepared biotin antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 μ L of prepared Streptavidin solution to each well.
7. Incubate 45 min at RT.
8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Assay Procedure: Prepare all reagents, samples and standards as instructed in the manual. Add 100 μ L of standard or sample to each well. Incubate 2.5 h at RT or O/N at 4 °C. Add 100 μ L of prepared biotin antibody to each well. Incubate 1 h at RT. Add 100 μ L of prepared Streptavidin solution to

Application Details

each well. Incubate 45 min at RT. Add 100 µL of TMB One-Step Substrate Reagent to each well. Incubate 30 min at RT. Add 50 µL of Stop Solution to each well. Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.

Expiry Date: 6 months
