

Datasheet for ABIN2748490

p53 ELISA Kit

[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	p53 (TP53)
Binding Specificity:	pSer15, total
Reactivity:	Human, Rat, Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human/Mouse/Rat Phospho-P53 (S15) and Total P53 ELISA Kit. This assay semi-quantitatively measures phosphorylated P53 (Ser15) and Total P53 in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes human Phospho-P53 (pSer15) and total (pan) P53.
Characteristics:	• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose) • Screen numerous different cell lysates without performing a Western Blot analysis • Minimal hands-on time, convenient, and non-radioactive material
Components:	• Pre-Coated 96-well Strip Microplate • Wash Buffer

Product Details

- Anti-Phospho Antibody
- Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- Streptavidin-Conjugated HRP
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:	• Distilled or deionized water • 100 mL and 1 liter graduated cylinders • Tubes to prepare sample dilutions • Protease and Phosphatase inhibitors • Precision pipettes to deliver 2 µL to 1 mL volumes • Adjustable 1-25 mL pipettes for reagent preparation • Benchtop rocker or shaker • Microplate reader capable of measuring absorbance at 450 nm
------------------------	--

Target Details

Target:	p53 (TP53)
Alternative Name:	P53 (TP53 Products)
Background:	P53 (Tumor protein p53, cellular tumor antigen p53, phosphoprotein p53, tumor suppressor p53) phosphorylated at Serine-15
Gene ID:	7157
UniProt:	P04637
Pathways:	p53 Signaling , MAPK Signaling , PI3K-Akt Signaling , Apoptosis , AMPK Signaling , Chromatin Binding , ER-Nucleus Signaling , Positive Regulation of Endopeptidase Activity , Hepatitis C , Protein targeting to Nucleus , Autophagy , Warburg Effect

Application Details

Sample Volume:	100 µL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents and samples as instructed in the manual. 2. Add 100 µL of sample or positive control to each well.

Application Details

- 3. Incubate 2.5 h at RT or O/N at 4 °C.
- 4. Add 100 µL of prepared primary antibody to each well.
- 5. Incubate 1 h at RT.
- 6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
- 7. Incubate 1 h at RT.
- 8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Assay Procedure: Prepare all reagents and samples as instructed in the manual.

Add 100 µL of sample or positive control to each well.

Incubate 2.5 h at RT or O/N at 4 °C.

Add 100 µL of prepared primary antibody to each well.

Incubate 1 h at RT.

Add 100 µL of prepared 1X HRP-Streptavidin to each well.

Incubate 1 h at RT.

Add 100 µL of TMB One-Step Substrate Reagent to each well.

Incubate 30 min at RT.

Add 50 µL of Stop Solution to each well.

Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months