antibodies -online.com





Datasheet for ABIN2748497

RPS6KB1 ELISA Kit



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	96 tests	
Target:	RPS6KB1	
Binding Specificity:	pSer424, pThr421	
Reactivity:	Human, Mouse, Rat	
Method Type:	Sandwich ELISA	
Application:	ELISA	
Product Details		
Purpose:	Human/Mouse/Rat Phospho-P70S6K (T421/S424) ELISA Kit. This assay semi-quantitatively measures phophorylated P70S6K (Thr421/S424) in lysate samples.	
Sample Type:	Cell Lysate, Tissue Lysate	
Analytical Method:	Semi-Quantitative	
Detection Method:	Colorimetric	
Specificity:	The antibody pair provided in this kit recognizes human, mouse, and rat p70S6 Kinase phosphorylated at sites Threonine-421 and/or Serine-424.	
Characteristics:	 Rapidly measure phosphorylated protein in lysates Screen numerous different cell lysates without performing a Western Blot analysis Minimal hands-on time, convenient, and non-radioactive material 	
Components:	Pre-Coated 96-well Strip MicroplateWash BufferAnti-Phospho Antibody	

Product Details

- · HRP-Conjugated Secondary Antibody
- · Assay Diluent
- · TMB One-Step Substrate
- · Stop Solution
- · Lysis Buffer
- · Positive Control Sample

Material not included:

- · Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- · Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	RPS6KB1
Alternative Name:	P70S6K (RPS6KB1 Products)
Background:	P70S6 kinase phosphorylated at T421 and/or S424
Gene ID:	6198
UniProt:	P23443
Pathways:	PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, Regulation of Cell Size, Skeletal Muscle
	Fiber Development, Feeding Behaviour, G-protein mediated Events, Smooth Muscle Cell
	Migration, Interaction of EGFR with phospholipase C-gamma, Warburg Effect

Application Details

Sample Volume:	100 μL	
Plate:	Pre-coated	
Protocol:	1. Prepare all reagents and samples as instructed in the manual.	
	2. Add 100 μ L of sample or positive control to each well.	
	3. Incubate 2.5 h at RT or O/N at 4 °C.	
	4. Add 100 μL of prepared primary antibody to each well.	
	5. Incubate 1 h at RT.	
	6. Add 100 μL of prepared 1X HRP-Streptavidin to each well.	

	7. Incubate 1 h at RT.	
	8. Add 100 μL of TMB One-Step Substrate Reagent to each well.	
	9. Incubate 30 min at RT.	
	10. Add 50 μL of Stop Solution to each well.11. Read at 450 nm immediately.	
Assay Procedure:	Prepare all reagents and samples as instructed in the manual.	
	Add 100 µL of sample or positive control to each well.	
	Incubate 2.5 h at RT or O/N at 4 °C.	
	Add 100 µL of prepared primary antibody to each well.	
	Incubate 1 h at RT.	
	Add 100 µL of prepared 1X HRP-Streptavidin to each well.	
	Incubate 1 h at RT.	
	Add 100 µL of TMB One-Step Substrate Reagent to each well.	
	Incubate 30 min at RT.	
	Add 50 µL of Stop Solution to each well.	
	Read at 450 nm immediately.	
Restrictions:	For Research Use only	
Handling		
Storage:	-20 °C	
Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of	
	shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-	
	Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell	
	Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return	
	unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20	
	°C. Reconstituted Positive Control (Item K) should be stored at -70 °C.	
Expiry Date:	6 months	