# antibodies .- online.com





# Datasheet for ABIN2748553

## **PF4 ELISA Kit**



## Overview

Quantity:	96 tests
Target:	PF4
Reactivity:	Rhesus Monkey
Method Type:	Sandwich ELISA
Detection Range:	0.041-10 ng/mL
Minimum Detection Limit:	0.041 ng/mL
Application:	ELISA
Product Details	
Purpose:	Rhesus Monkey Platelet Factor 4 (CXCL4) ELISA Kit for serum, plasma, and cell culture
	supernatant samples.
Sample Type:	Plasma, Cell Culture Supernatant, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA antibody pair detects Rhesus Macaque PF-4 and human PF-4. Other species not
	determined
Characteristics:	Strip plates and additional reagents allow for use in multiple experiments
	Quantitative protein detection
	Establishes normal range  The bound of the first in the second of t
	The best products for confirmation of antibody array data

## **Product Details**

#### Components:

- · Pre-Coated 96-well Strip Microplate
- Wash Buffer
- · Stop Solution
- Assay Diluent(s)
- · Lyophilized Standard
- · Biotinylated Detection Antibody
- · Streptavidin-Conjugated HRP
- TMB One-Step Substrate

#### Material not included:

- Distilled or deionized water
- Precision pipettes to deliver 2 μL to 1 μL volumes
- Adjustable 1-25 µL pipettes for reagent preparation
- 100 µL and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

# Target Details

Target:	PF4
Alternative Name:	Platelet Factor 4 (PF4 Products)
Background:	Platelet Factor 4 (CXCL4)
Gene ID:	5196
UniProt:	P02776

#### **Application Details**

Application Notes:	Recommended Dilution for serum and plasma samples5,000 fold
Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.
	2. Add 100 $\mu L$ of standard or sample to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 $\mu L$ of prepared biotin antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 $\mu L$ of prepared Streptavidin solution to each well.
	7. Incubate 45 min at RT.

# **Application Details**

	<ul><li>8. Add 100 μL of TMB One-Step Substrate Reagent to each well.</li><li>9. Incubate 30 min at RT.</li></ul>
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Assay Procedure:	Prepare all reagents, samples and standards as instructed in the manual.Add 100 μL of
	standard or sample to each well.Incubate 2.5 h at RT or O/N at 4 °C.Add 100 µL of prepared
	biotin antibody to each well. Incubate 1 h at RT. Add 100 µL of prepared Streptavidin solution to
	each well.Incubate 45 min at RT.Add 100 μL of TMB One-Step Substrate Reagent to each
	well.Incubate 30 min at RT.Add 50 μL of Stop Solution to each well.Read at 450 nm
	immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated
	freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is
	recommended to store at -80°C.
Expiry Date:	6 months