

Datasheet for ABIN2748634 **CXCL12 ELISA Kit**

[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	CXCL12
Reactivity:	Rhesus Monkey
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Custom Rhesus Macaque Stromal Cell-Derived Factor 1 (SDF1) ELISA Kit.
Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Rhesus Macaque Stromal Cell-Derived Factor 1 (SDF1)
Characteristics:	• Strip plates and additional reagents allow for use in multiple experiments • Quantitative protein detection • Establishes normal range • The best products for confirmation of antibody array data
Components:	• Pre-Coated 96-well Strip Microplate • Wash Buffer • Stop Solution • Assay Diluent(s) • Lyophilized Standard

Product Details

- Biotinylated Detection Antibody
- Streptavidin-Conjugated HRP
- TMB One-Step Substrate

Material not included:	• Distilled or deionized water • Precision pipettes to deliver 2 µL to 1 µL volumes • Adjustable 1-25 µL pipettes for reagent preparation • 100 µL and 1 liter graduated cylinders • Tubes to prepare standard and sample dilutions • Absorbent paper • Microplate reader capable of measuring absorbance at 450nm • Log-log graph paper or computer and software for ELISA data analysis
------------------------	---

Target Details

Target:	CXCL12
Alternative Name:	SDF1 (CXCL12 Products)
Background:	Stromal Cell-Derived Factor 1 (SDF1)
Gene ID:	574334
UniProt:	Q8HYP0
Pathways:	Regulation of Cell Size , CXCR4-mediated Signaling Events , Negative Regulation of intrinsic apoptotic Signaling

Application Details

Sample Volume:	100 µL
Plate:	Pre-coated
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual. 2. Add 100 µL of standard or sample to each well. 3. Incubate 2.5 h at RT or O/N at 4 °C. 4. Add 100 µL of prepared biotin antibody to each well. 5. Incubate 1 h at RT. 6. Add 100 µL of prepared Streptavidin solution to each well. 7. Incubate 45 min at RT. 8. Add 100 µL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 µL of Stop Solution to each well. 11. Read at 450 nm immediately.

Application Details

Assay Procedure: Prepare all reagents, samples and standards as instructed in the manual. Add 100 µL of standard or sample to each well. Incubate 2.5 h at RT or O/N at 4 °C. Add 100 µL of prepared biotin antibody to each well. Incubate 1 h at RT. Add 100 µL of prepared Streptavidin solution to each well. Incubate 45 min at RT. Add 100 µL of TMB One-Step Substrate Reagent to each well. Incubate 30 min at RT. Add 50 µL of Stop Solution to each well. Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: The entire kit may be stored at -20°C for up to 1 year from the date of shipment. Avoid repeated freeze-thaw cycles. The kit may be stored at 4°C for up to 6 months. For extended storage, it is recommended to store at -80°C.

Expiry Date: 6 months