

Datasheet for ABIN2749108  
**anti-SLAMF1 antibody (FITC)**



[Go to Product page](#)

2 Images

2 Publications

## Overview

Quantity:	100 tests
Target:	SLAMF1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SLAMF1 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

## Product Details

Immunogen:	Human CD150-transfected 300.19 cells
Clone:	SLAM-4
Isotype:	IgG1
Specificity:	The mouse monoclonal antibody SLAM.4 recognizes an extracellular epitope of CD150, a cell surface molecule expressed on lymphocytes and involved in their activation.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

## Target Details

Target:	SLAMF1
---------	--------

## Target Details

Alternative Name:	CD150 ( <a href="#">SLAMF1 Products</a> )
Background:	Signaling lymphocytic activation molecule family m,CD150, also known as SLAM (signaling lymphocyte activation molecule) is a 70-95 kDa single chain transmembrane phosphoglycoprotein of the CD2 family. Its extracellular part contains eight potential N-glycosylation sites, and the intracellular tail contains three unique tyrosine-based motifs. These binding sites can be recognized by SH2-binding phosphatases and the adaptor proteins, such as SAP/SH2D1A or EAT-2. The SLAM family receptors are involved in leucocyte activation and contribute to the effective germinal center formation, generation of high-affinity antibody-secreting plasma cells, and memory T and B cells, thereby facilitating long-term immune response. CD150 expression is upregulated after cell activation.,SLAM, SLAMF1
Gene ID:	6504
UniProt:	<a href="#">Q13291</a>

## Application Details


Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µL reagent / 100 µL of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.
Comment:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Restrictions:	For Research Use only

## Handling

Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.


Product cited in: Romero, Zapater, Calvo, Kalko, de la Fuente, Tovar, Ockeloen, Pizcueta, Engel: "CD229 (Ly9) lymphocyte cell surface receptor interacts homophilically through its N-terminal domain and relocalizes to the immunological synapse." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 174, Issue 11, pp. 7033-42, (2005) ([PubMed](#)).

Romero, Benítez, March, Vilella, Miralpeix, Engel: "Differential expression of SAP and EAT-2-binding leukocyte cell-surface molecules CD84, CD150 (SLAM), CD229 (Ly9) and CD244 (2B4)." in: **Tissue antigens**, Vol. 64, Issue 2, pp. 132-44, (2004) ([PubMed](#)).



**Flow Cytometry**

**Image 1.** Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD150 (SLAM.4) FITC antibody (4 µL reagent / 100 µL of peripheral whole blood).



**Flow Cytometry**

**Image 2.** Separation of human CD150 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD150 (SLAM.4) FITC antibody (4 µL reagent / 100 µL of peripheral whole blood).