

Datasheet for ABIN2749113
anti-CD180 antibody (APC)



[Go to Product page](#)

1 Image 3 Publications

Overview

Quantity:	100 tests
Target:	CD180
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD180 antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

Product Details

Immunogen:	Human tonsillar B cells
Clone:	G28-8
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody G28-8 recognizes an extracellular epitope of CD180, a 95-105 kDa TLR-like glycoprotein expressed on peripheral blood monocytes and dendritic cells, mantle zone B cells and marginal zone B cells, but very weakly on germinal center B cells.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

Target:	CD180
Alternative Name:	CD180 (CD180 Products)
Background:	CD180 Molecule,CD180, also known as RP105 (or Bgp95, LY64) is a type I membrane glycoprotein of Toll-like receptor (TLR) family. Its cytoplasmic tail is short and unlike the TLRs, it lacks the TIR domain. CD180 expression is dependent on the coexpression of its helper molecule, MD-1, and mirrors that of TLR4 on antigen-presenting cells. CD180 regulates recognition of LPS and signaling in B cells, via interacting directly with the TLR4 signaling complex, inhibiting its ability to bind microbial ligands. Ligation of CD180 by monoclonal antibodies leads to B cell activation, upregulation of CD80/CD86, and increase in cell size.,LY64, Ly78, RP105, Bgp95
Gene ID:	4064
UniProt:	Q99467
Pathways:	Cellular Response to Molecule of Bacterial Origin , Toll-Like Receptors Cascades

Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µL reagent / 100 µL of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Comment:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Restrictions:	For Research Use only

Handling

Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.


Publications

Product cited in: Olson, Graves, Shu, Ryan, Clark: "Caspase activity is required for stimulated B lymphocytes to enter the cell cycle." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 170, Issue 12, pp. 6065-72, (2003) ([PubMed](#)).

Clark, Shu, Lüscher, Draves, Banchereau, Ledbetter, Valentine: "Activation of human B cells. Comparison of the signal transduced by IL-4 to four different competence signals." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 143, Issue 12, pp. 3873-80, (1990) ([PubMed](#)).

Valentine, Clark, Shu, Norris, Ledbetter: "Antibody to a novel 95-kDa surface glycoprotein on human B cells induces calcium mobilization and B cell activation." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 140, Issue 12, pp. 4071-8, (1988) ([PubMed](#)).

Images



Flow Cytometry

Image 1. Flow cytometry analysis (surface staining) of human peripheral blood cells with anti-CD180 (G28-8) APC.