antibodies -online.com





anti-CD94 antibody (APC)





Publications



Go to Product page

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	100 tests	
Target:	CD94 (KLRD1)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CD94 antibody is conjugated to APC	
Application:	Flow Cytometry (FACS)	

Product Details

Immunogen:	Cultured human NK cells
Clone:	HP-3D9
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody HP-3D9 recognizes an extracellular epitope of CD94, a 70 kDa type II transmembrane glycoprotein expressed on NK cells, NK-T cells, and subsets of CD8+ T cells and gamma/delta T cells.
Cross-Reactivity (Details):	Human
Purification:	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

rargerbetane		
Target:	CD94 (KLRD1)	
Alternative Name:	CD94 (KLRD1 Products)	
Background:	Killer cell lectin like receptor D1,CD94, also known as KLRD1 (killer cell lectin-like receptor D1), is a transmembrane glycoprotein of the C-type lectin family, which forms disulfide-linked heterodimers with NKG2A, B, C, E, H proteins, constituting functionally distinct receptors of NK cells and related cell types. CD94/NKG2A and CD94/NKG2B heterodimers serve as inhibitory, whereas CD94/NKG2C and CD94/NKG2E as activating receptors. The ligand for CD94/NKG2 complexes has been identified as HLA-E. Extent of CD94 expression on NK cell surface can be used to demonstrate their progress through the differentiation process.,KLR-D1, KLRD1, KCLLRD1	
Gene ID:	3824	
UniProt:	Q13241	
Application Details		
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.	
Comment:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.	
Restrictions:	For Research Use only	
Handling		
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	

Product cited in:

Yu, Mao, Wei, Hughes, Zhang, Park, Liu, McClory, Marcucci, Trotta, Caligiuri: "CD94 surface density identifies a functional intermediary between the CD56bright and CD56dim human NK-cell subsets." in: **Blood**, Vol. 115, Issue 2, pp. 274-81, (2010) (PubMed).


Bovenschen, Van De Kerkhof, Gerritsen, Seyger: "The role of lesional T cells in recalcitrant psoriasis during infliximab therapy." in: **European journal of dermatology: EJD**, Vol. 15, Issue 6, pp. 454-8, (2005) (PubMed).

Wada, Matsumoto, Maenaka, Suzuki, Yamamoto: "The inhibitory NK cell receptor CD94/NKG2A and the activating receptor CD94/NKG2C bind the top of HLA-E through mostly shared but partly distinct sets of HLA-E residues." in: **European journal of immunology**, Vol. 34, Issue 1, pp. 81-90, (2004) (PubMed).

Ntrivalas, Kwak-Kim, Gilman-Sachs, Chung-Bang, Ng, Beaman, Mantouvalos, Beer: "Status of peripheral blood natural killer cells in women with recurrent spontaneous abortions and infertility of unknown aetiology." in: **Human reproduction (Oxford, England)**, Vol. 16, Issue 5, pp. 855-61, (2001) (PubMed).

Seo, Tokura, Ishihara, Takeoka, Tagawa, Takigawa: "Disordered expression of inhibitory receptors on the NK1-type natural killer (NK) leukaemic cells from patients with hypersensitivity to mosquito bites." in: **Clinical and experimental immunology**, Vol. 120, Issue 3, pp. 413-9, (2000) (PubMed).

There are more publications referencing this product on: Product page



Flow Cytometry

Image 1. Surface staining of CD94 in human peripheral blood with anti-CD94 (HP-3D9) APC.