-online.com antibodies

Datasheet for ABIN2754657 anti-hCoV-OC43 Spike antibody (AA 15-344) (Biotin)



Overview

| Quantity: | 100 µg |
|----------------------|---|
| Target: | hCoV-0C43 Spike (HCoV-0C43 S) |
| Binding Specificity: | AA 15-344 |
| Reactivity: | Human Coronavirus OC43 (HCoV-OC43) |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This hCoV-OC43 Spike antibody is conjugated to Biotin |
| Application: | ELISA |
| | |
| Product Details | |

| Immunogen: | Recombinant Human coronavirus OC43 Spike glycoprotein protein(15-344aa) |
|------------------------------|---|
| lsotype: | lgG |
| Purification: | Caprylic Acid Ammonium Sulfate Precipitation |
| T () () | |
| Target Details | |
| | |
| Target: | hCoV-0C43 Spike (HCoV-0C43 S) |
| Target: Alternative Name: | hCoV-0C43 Spike (HCoV-0C43 S) 0C43 Spike Glycoprotein (HCoV-0C43 S Products) |
| | |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2754657 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

| Target Details | |
|---------------------|---|
| | protein has at least 3 conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and target cell membrane fusion, the coiled coil regions (heptad repeats) assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of viral and target cell membranes. |
| UniProt: | P36334 |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4 |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C/-20 °C/-80 °C |