

## Datasheet for ABIN2773953

# anti-UXS1 antibody (Middle Region)



Overview



Go to Product page

Quantity:	100 μL	
Target:	UXS1	
Binding Specificity:	Middle Region	
Reactivity:	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Zebrafish (Danio rerio)	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UXS1 antibody is un-conjugated	

### Application: Western Blotting (WB)

#### **Product Details**

Immunogen:	The immunogen is a synthetic peptide directed towards the middle region of human UXS1
Sequence:	LMLGWEPVVP LEEGLNKAIH YFRKELEYQA NNQYIPKPKP ARIKKGRTRH
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rat: 100%, Zebrafish: 93%
Characteristics:	This is a rabbit polyclonal antibody against UXS1. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

## Target Details

Target: UXS1


## Target Details

Alternative Name:	UXS1 (UXS1 Products)
Background:	UDP-glucuronate decarboxylase (UGD, EC 4.1.1.35) catalyzes the formation of UDP-xylose from
	UDP-glucuronate. UDP-xylose is then used to initiate glycosaminoglycan biosynthesis on the
	core protein of proteoglycans.UDP-glucuronate decarboxylase (UGD, EC 4.1.1.35) catalyzes the
	formation of UDP-xylose from UDP-glucuronate. UDP-xylose is then used to initiate
	glycosaminoglycan biosynthesis on the core protein of proteoglycans.[supplied by OMIM].
	PRIMARYREFSEQ_SPAN PRIMARY_IDENTIFIER PRIMARY_SPAN COMP 1-49 BI825770.1 1-49
	50-2073 AY358541.1 1-2024 2074-2132 BM472113.1 164-222
	Alias Symbols: FLJ23591, UGD, SDR6E1
	Protein Interaction Partner: AKT1,
	Protein Size: 420
Molecular Weight:	47 kDa
Gene ID:	80146
NCBI Accession:	NM_025076, NP_079352
UniProt:	Q8NBZ7
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 420 AA
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

#### Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Images**



#### **Western Blotting**

**Image 1.** WB Suggested Anti-UXS1 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:312500 Positive Control: OVCAR-3 cell lysate UXS1 is supported by BioGPS gene expression data to be expressed in OVCAR3