antibodies .- online.com







anti-ZNF771 antibody (N-Term)





Publication



()	1 /	-	r٧	/ 1	0	A .
	1//	\vdash	1 \/	/ I	_	۱/۱
\sim	v	\sim	1 V		$\overline{}$	V١

Quantity:	100 μL
Target:	ZNF771
Binding Specificity:	N-Term
Reactivity:	Human, Guinea Pig, Cow, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF771 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human ZNF771
Sequence:	EVVKLKIPMD NKEVPGEAPA PSADPARPHA CPDCGRAFAR RSTLAKHART
Predicted Reactivity:	Cow: 93%, Dog: 86%, Guinea Pig: 79%, Human: 100%
Characteristics:	This is a rabbit polyclonal antibody against ZNF771. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified
Target Details	
Target:	ZNF771

Target Details

Alternative Name:	ZNF771 (ZNF771 Products)
Background:	The specific function of this protein remains unknown.
	Alias Symbols: DSC43
	Protein Interaction Partner: SUMO1,
	Protein Size: 275
Molecular Weight:	31 kDa
Gene ID:	51333
NCBI Accession:	NM_016643, NP_057727
UniProt:	B2R9V3

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 275 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.


Publications

Product cited in: Lazrek, Goffard, Schanen, Karquel, Bocket, Lion, Devaux, Hedouin, Gosset, Hober: "Detection of

hepatitis C virus antibodies and RNA among medicolegal autopsy cases in Northern France." in:

Diagnostic microbiology and infectious disease, Vol. 55, Issue 1, pp. 55-8, (2006) (PubMed).

Images



Western Blotting

Image 1. WB Suggested Anti-ZNF771 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:1562500 Positive Control: Human Thymus