# antibodies .- online.com





## anti-ASCC2 antibody (Middle Region)



Image



Go to Product page

$\sim$			
	N/6	1//r	$I \cap V$


Quantity:	100 μL	
Target:	ASCC2	
Binding Specificity:	Middle Region	
Reactivity:	Human, Mouse, Rat, Rabbit, Cow, Dog, Guinea Pig, Horse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ASCC2 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	The immunogen is a synthetic peptide directed towards the middle region of human ASCC2	
Immunogen: Sequence:	The immunogen is a synthetic peptide directed towards the middle region of human ASCC2 YEDEYDDTYD GNQVGANDAD SDDELISRRP FTIPQVLRTK VPREGQEEDD	
Sequence:	YEDEYDDTYD GNQVGANDAD SDDELISRRP FTIPQVLRTK VPREGQEEDD  Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit:	
Sequence: Predicted Reactivity:	YEDEYDDTYD GNQVGANDAD SDDELISRRP FTIPQVLRTK VPREGQEEDD  Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%  This is a rabbit polyclonal antibody against ASCC2. It was validated on Western Blot using a cell	
Sequence:  Predicted Reactivity:  Characteristics:	YEDEYDDTYD GNQVGANDAD SDDELISRRP FTIPQVLRTK VPREGQEEDD  Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%  This is a rabbit polyclonal antibody against ASCC2. It was validated on Western Blot using a cell lysate as a positive control.	

## **Target Details**

Alternative Name:	ASCC2 (ASCC2 Products)		
Background:	ASCC2 belongs to the ASCC2 family. It contains 1 CUE domain. ASCC2 enhances NF-kappa-B,		
	SRF and AP1 transactivation.		
	Alias Symbols: ASC1p100, p100		
	Protein Interaction Partner: UBC, GADD45A, RNF11, PCBD2, POLDIP2, CKAP4, TBC1D17, OLA1		
	MED31, CUTC, URM1, POLA2, IGSF9, KDM1A, RADIL, GTF3C1, FUNDC2, GNL3, ELAC2, TRIP4,		
	FAF1, ASS1, SNW1, PIK3CD, DEAF1, GFER, IMMT, PJA1, PIN4, LPL, FBP1, SNRPB, RPLP1,		
	RPA1, MYH9, EEF1D, SRF, RELA, J		
	Protein Size: 757		
Molecular Weight:	86 kDa		
Gene ID:	84164		
NCBI Accession:	NM_032204, NP_115580		
UniProt:	Q9H1I8		
Application Details			
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.		
Comment:	Antigen size: 757 AA		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	Lot specific		
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 %		
	sucrose.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
Handling Advice:	Avoid repeated freeze-thaw cycles.		
Storage:	-20 °C		
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small		

aliquots to prevent freeze-thaw cycles.

### **Images**



### **Western Blotting**

**Image 1.** WB Suggested Anti-ASCC2 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:12500 Positive Control: HepG2 cell lysate ASCC2 is supported by BioGPS gene expression data to be expressed in HepG2