

Datasheet for ABIN2776851 anti-MUC1 antibody (N-Term)





Go to Floudet page

Overview	
Quantity:	100 μL
Target:	MUC1
Binding Specificity:	N-Term
Reactivity:	Human, Cow, Dog, Pig, Rabbit, Goat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MUC1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human MUC1
Sequence:	SATQRSSVPS STEKNALSTG VSFFFLSFHI SNLQFNSSLE DPSTDYYQEL
Predicted Reactivity:	Cow: 86%, Dog: 86%, Goat: 86%, Human: 100%, Pig: 82%, Rabbit: 86%
Characteristics:	This is a rabbit polyclonal antibody against MUC1. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified
Target Details	
Target:	MUC1
Alternative Name:	MUC1 (MUC1 Products)

Background:

MUC1 is a membrane bound, glycosylated phosphoprotein. The protein is anchored to the apical surface of many epithelia by a transmembrane domain, with the degree of glycosylation varying with cell type. It also includes a 20 aa variable number tandem repeat (VNTR) domain, with the number of repeats varying from 20 to 120 in different individuals. The protein serves a protective function by binding to pathogens and also functions in a cell signaling capacity. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is a member of the mucin family and encodes a membrane bound, glycosylated phosphoprotein. The protein is anchored to the apical surface of many epithelia by a transmembrane domain, with the degree of glycosylation varying with cell type. It also includes a 20 aa variable number tandem repeat (VNTR) domain, with the number of repeats varying from 20 to 120 in different individuals. The protein serves a protective function by binding to pathogens and also functions in a cell signaling capacity. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. Multiple alternatively spliced transcript variants that encode different isoforms of this gene have been reported, but the full-length nature of only some has been determined.

Alias Symbols: CD227, EMA, H23AG, MAM6, PEM, PEMT, PUM, KL-6, MUC-1, MUC-1/X, MUC1/ZD, MUC-1/SEC

Protein Interaction Partner: UBC, PPARG, IPO5, NUP62, EGFR, ABL1, SRC, JUN, TP53, TBP, IKBKG, TNFRSF1A, IKBKB, GRB2, ESR1, GALNT15, GALNT12, GALNT10, OSGEP, GSK3B, GALNT2, GALNT1, CTNND1, SIGLEC1, ERBB4, ERBB3, SOS1, PRKCD, ZAP70, APC, JUP, LYN, ERBB2, LCK, CTNNB1, GALNT4, BAX, CDKN1

Protein Size: 273

Molecular Weight:	27 kDa
Gene ID:	4582
NCBI Accession:	NM_002456, NP_002447
UniProt:	P15941
Pathways:	Negative Regulation of intrinsic apoptotic Signaling


Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 273 AA
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images




Western Blotting

Image 1.



Immunohistochemistry

Image 2.



Western Blotting

Image 3. Host: Rabbit Target Name: MUC1 Sample Type: 721_B Antibody Dilution: 1.0ug/ml There is BioGPS gene expression data showing that MUC1 is expressed in 721_B