# antibodies - online.com







# anti-Cyclin H antibody (N-Term)



**Images** 



$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL	
Target:	Cyclin H (CCNH)	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat, Guinea Pig, Rabbit, Horse, Dog, Cow	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Cyclin H antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human CCNH	
Sequence:	PHEEMTLCKY YEKRLLEFCS VFKPAMPRSV VGTACMYFKR FYLNNSVMEY	
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%	

### **Target Details**

Characteristics:

Purification:

Target:	Cyclin H (CCNH)
Alternative Name:	CCNH (CCNH Products)

Affinity Purified

This is a rabbit polyclonal antibody against CCNH. It was validated on Western Blot.


## **Target Details**

Background:	Cyclin H regulates CDK7, the catalytic subunit of the CDK- activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDC2/CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIH basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive carboxyl-terminus domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. It is involved in cell cycle control and in RNA transcription by RNA polymerase II. Its expression and activity are constant throughout the cell cycle. Alias Symbols: CAK, p34, p37  Protein Interaction Partner: SFN, GOLGA2, SSX2IP, CCDC170, CCDC33, DUSP12, SSSCA1,	
	NDC80, CALCOCO2, PPFIA1, PSMA1, KLC3, RPA3, RPA2, RPA1, NEDD4, CDK7, CDK8, CDK2, CCNC, MNAT1, MBP, CTD, CDK20, CTBP2, CSNK2B, TRIM8, GANAB, RARB, POLR2A, GTF2H4, ERCC2, tat, POLR2B, CDK9, BRCA1, GTF2H1  Protein Size: 323	
Molecular Weight:	38 kDa	
Gene ID:	902	
NCBI Accession:	NM_001239, NP_001230	
UniProt:	P51946	
Pathways:	Cell Division Cycle, Mitotic G1-G1/S Phases, M Phase	
Application Details		
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.	
Comment:	Antigen size: 323 AA	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	

#### Handling

	should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Images**



31 kDa\_\_

22 kDa

#### **Western Blotting**

**Image 1.** WB Suggested Anti-CCNH Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:62500 Positive Control: Hela cell lysate CCNH is supported by BioGPS gene expression data to be expressed in HeLa


#### **Western Blotting**

Image 2. WB Suggested Anti-CCNH

Antibody Titration: 0.2-1  $\mu$ g/mL ELISA Titer: 1:.2500

Positive Control: Hela cell lysate

CCNH is supported by BioGPS gene expression data to be expressed in HeLa



#### **Western Blotting**

**Image 3.** Host: Mouse Target Name: CCNH Sample Tissue: Mouse Skeletal Muscle Antibody Dilution: 1ug/ml