

Datasheet for ABIN2778019
anti-ZNF768 antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	ZNF768
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Rabbit, Cow, Horse, Dog, Guinea Pig, Saccharomyces cerevisiae
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF768 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human FLJ23436
Sequence:	GLEPQSPGFE PQSPEFEPQS PRFEPESPGF ESRSPGLVPP SPEFAPRSPE
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Yeast: 100%
Characteristics:	This is a rabbit polyclonal antibody against FLJ23436. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Protein A purified

Target Details

Target:	ZNF768
---------	--------

Target Details

Abstract:	ZNF768 Products
Background:	FLJ23436 is a hypothetical protein found in Chromosome 16 Protein Interaction Partner: PASK, MDC1, BARD1, UBC, TNNT1, NDUFA12, HMP19, ZNF277, Protein Size: 520
Molecular Weight:	58 kDa
Gene ID:	79724
NCBI Accession:	NM_024671 , NP_078947
UniProt:	Q9H5H4

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 520 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western Blotting

Image 1. WB Suggested Anti-FLJ23436 Antibody Titration:
1.25ug/ml Positive Control: HepG2 cell lysate