antibodies - online.com







anti-ZNF567 antibody (N-Term)





Publication



()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

OVEIVIEW	
Quantity:	100 μL
Target:	ZNF567 (104092)
Binding Specificity:	N-Term
Reactivity:	Human, Cow, Dog, Horse, Pig, Rabbit, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF567 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human ZNF567
Sequence:	TSFAGHTCLE ENWKAEDFLV KFKEHQEKYS RSVVSINHKK LVKEKSKIYE
Predicted Reactivity:	Cow: 93%, Dog: 93%, Horse: 100%, Human: 100%, Pig: 100%, Rabbit: 93%, Rat: 92%
Characteristics:	This is a rabbit polyclonal antibody against ZNF567. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified
. d.moddori.	running i dinied
Target Details	
Target:	ZNF567 (104092)

Target Details

Alternative Name:	ZNF567 (104092 Products)
Background:	ZNF567 may be involved in transcriptional regulation. Alias Symbols: MGC45586 Protein Interaction Partner: FAM9B, ZAK, UBC, PCBD1, Protein Size: 616
Molecular Weight:	71 kDa
Gene ID:	163081
NCBI Accession:	NM_152603, NP_689816

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.	
Comment:	Antigen size: 616 AA	
Restrictions:	For Research Use only	

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Publications


Product cited in:

Wang, Mueller, Hertel, Cambi: "G Run-mediated recognition of proteolipid protein and DM20 5" splice sites by U1 small nuclear RNA is regulated by context and proximity to the splice site." in:

The Journal of biological chemistry, Vol. 286, Issue 6, pp. 4059-71, (2011) (PubMed).

Wang, Cambi: "Heterogeneous nuclear ribonucleoproteins H and F regulate the proteolipid protein/DM20 ratio by recruiting U1 small nuclear ribonucleoprotein through a complex array of G runs." in: **The Journal of biological chemistry**, Vol. 284, Issue 17, pp. 11194-204, (2009) (PubMed).

Images



Western Blotting

Image 1. WB Suggested Anti-ZNF567 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:1562500 Positive Control: PANC1 cell lysate