antibodies -online.com







Images



Go to Product page

Overview

| Quantity: | 100 μL |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| Target: | AIF (AIFM1) |
| Binding Specificity: | C-Term |
| Reactivity: | Human, Mouse, Rat, Pig, Cow, Dog, Horse, Sheep, Rabbit, Zebrafish (Danio rerio), Guinea Pig |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AIF antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC) |
| Product Details | |
| Immunogen: | The immunogen is a synthetic peptide directed towards the C terminal region of human PDCD8 |
| Sequence: | VDSSLPTVGV FAKATAQDNP KSATEQSGTG IRSESETESE ASEITIPPST |
| Predicted Reactivity: | Cow: 100%, Dog: 100%, Guinea Pig: 93%, Horse: 100%, Human: 100%, Mouse: 100%, Pig: 100%, Rabbit: 93%, Rat: 100%, Sheep: 100%, Zebrafish: 79% |
| Characteristics: | This is a rabbit polyclonal antibody against PDCD8. It was validated on Western Blot using a cell lysate as a positive control. |
| Purification: | Affinity Purified |
| Target Details | |
| Target: | AIF (AIFM1) |
| | |


Target Details


| Alternative Name: | PDCD8 (AIFM1 Products) |
|---------------------|---------------------------------------------------------------------------------------------------|
| Background: | AIFM1 (PDCD8) is a flavoprotein essential for nuclear disassembly in apoptotic cells that is |
| | found in the mitochondrial intermembrane space in healthy cells. Induction of apoptosis result |
| | in the translocation of this protein to the nucleus where it effects chromosome condensation |
| | and fragmentation. In addition, AIFM1 induces mitochondria to release the apoptogenic |
| | proteins cytochrome c and caspase-9. This gene encodes a flavoprotein essential for nuclear |
| | disassembly in apoptotic cells that is found in the mitochondrial intermembrane space in |
| | healthy cells. Induction of apoptosis results in the translocation of this protein to the nucleus |
| | where it effects chromosome condensation and fragmentation. In addition, this gene product |
| | induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. Three |
| | alternative transcripts encoding different isoforms have been identified for this gene. |
| | Alias Symbols: AIF, PDCD8, COXPD6 |
| | Protein Interaction Partner: ISG15, STAU1, BAG6, UBC, NEDD8, MDM2, ASB12, RNF2, BMI1, |
| | SUZ12, PARK2, BAG3, ADRB2, HDAC11, OXSR1, ILK, PAN2, NOS2, MLH1, CFTR, ESR1, TST, |
| | FSCN1, CPOX, SHC1, FBX06, TSC22D4, CAND1, COPS5, CUL1, CUL2, CUL3, CUL5, TONSL, |
| | BLNK, PIDD1, XIAP, AMFR, Tubb4b, Tub |
| | Protein Size: 326 |
| Molecular Weight: | 36 kDa |
| Gene ID: | 9131 |
| NCBI Accession: | NM_145813, NP_665812 |
| Pathways: | Apoptosis, Positive Regulation of Endopeptidase Activity, Cell RedoxHomeostasis, Smooth |
| | Muscle Cell Migration, Warburg Effect |
| Application Details | |
| Application Notes: | Optimal working dilutions should be determined experimentally by the investigator. |
| Comment: | Antigen size: 326 AA |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % |
| | |

Handling

| | sucrose. |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | -20 °C |
| Storage Comment: | For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. |

Images





Western Blotting

Image 1. WB Suggested Anti-PDCD8 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:62500 Positive Control: Jurkat cell lysate AIFM1 is supported by BioGPS gene expression data to be expressed in Jurkat


Western Blotting

Image 2. WB Suggested Anti-PDCD8

Antibody Titration: 0.2-1 µg/mL ELISA Titer: 1:.2500

Positive Control: Jurkat cell lysate

AIFM1 is supported by BioGPS gene expression data to be expressed in Jurkat



Western Blotting

Image 3. Host: Rabbit Target Name: AIFM1 Sample Tissue:
Mouse Liver Antibody Dilution: 1ug/ml

Please check the product details page for more images. Overall 5 images are available for ABIN2778665.