antibodies .- online.com



Datasheet for ABIN2779092

anti-Pre-mRNA Branch Site Protein p14 (SF3B14) (N-Term) antibody



Go to Product pag

Image	
irriage	

1

Publication

	Go to Froduct page
Overview	
Quantity:	100 μL
Target:	Pre-mRNA Branch Site Protein p14 (SF3B14)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Zebrafish (Danio rerio), Dog, Guinea Pig, Horse, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human SF3B14
Sequence:	MAMQAAKRAN IRLPPEVNRI LYIRNLPYKI TAEEMYDIFG KYGPIRQIRV
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 100%
Characteristics:	This is a rabbit polyclonal antibody against SF3B14. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target: Pre-mRNA Branch Site Protein p14 (SF3B14)


Target Details

Alternative Name:	SF3B14 (SF3B14 Products)
Background:	SF3B14 is a 14 kDa protein subunit of the splicing factor 3b complex. Splicing factor 3b
	associates with both the U2 and U11/U12 small nuclear ribonucleoprotein complexes (U2
	snRNP) of spliceosomes. This 14 kDa protein interacts directly with subunit 1 of the splicing
	factor 3b complex. SF3B14 also interacts directly with the adenosine that carries out the first
	transesterification step of splicing at the pre-mRNA branch site. This gene encodes a 14 kDa
	protein subunit of the splicing factor 3b complex. Splicing factor 3b associates with both the U.
	and U11/U12 small nuclear ribonucleoprotein complexes (U2 snRNP) of spliceosomes. This 14
	kDa protein interacts directly with subunit 1 of the splicing factor 3b complex. This 14 kDa
	protein also interacts directly with the adenosine that carries out the first transesterification
	step of splicing at the pre-mRNA branch site.
	Alias Symbols: CGI-110, HSPC175, Ht006, P14, SAP14, SF3B14a
	Protein Interaction Partner: UBC, WWOX, SUZ12, EED, PAN2, ITGA4, FN1, SF3B1, VCAM1,
	SF3B5, THOC6, WDR18, TRA2A, PRPF19, RBMX, SF3B3, PUF60, DHX15, RCC1, APP, U2AF2,
	ZNF326, ZC3H18, FTSJ3, PHF5A, TRIM55, SF3B2, UTP14A, USP39, PRPF8, BCAS2, SF3B4,
	THRAP3, EFTUD2, SRSF11, SART1, VTN, U2
	Protein Size: 125
Molecular Weight:	14 kDa
Gene ID:	51639
NCBI Accession:	NM_016047, NP_057131
UniProt:	Q9Y3B4
Application Details	
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 125 AA
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 %
	sucrose.

Handling

Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Publications	
Product cited in:	Huang, Ma, Li, Yu, Zhang, Wei, Jin, Xu, Gao, Huang: "NF-кВ1 inhibits c-Myc protein degradation

Images



PubMed).

Western Blotting

through suppression of FBW7 expression." in: Oncotarget, Vol. 5, Issue 2, pp. 493-505, (2014) (

Image 1. WB Suggested Anti-SF3B14 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:312500 Positive Control: THP-1 cell lysate