



[Go to Product page](#)

Datasheet for ABIN2779435

anti-NFATC4 antibody (N-Term)

2 Images

Overview

Quantity:	100 µL
Target:	NFATC4
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Dog, Cow, Horse, Rabbit, Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFATC4 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human NFATC4
Sequence:	RGPEDSWLLL SAPGTPASP RPASPCGKRR YSSSGTPSSA SPALSRRGSL
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%
Characteristics:	This is a rabbit polyclonal antibody against NFATC4. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target:	NFATC4
---------	--------

Target Details

Alternative Name: [NFATC4 \(NFATC4 Products\)](#)

Background: NFATC4 is a member of the nuclear factors of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family of nuclear factors of activated T cells also participate in the formation of this complex. NFATC4 plays a role in the inducible expression of cytokine genes in T cells, especially in the induction of the IL-2 and IL-4. The product of this gene is a member of the nuclear factors of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family of nuclear factors of activated T cells also participate in the formation of this complex. The product of this gene plays a role in the inducible expression of cytokine genes in T cells, especially in the induction of the IL-2 and IL-4.

Alias Symbols: NF-ATc4, NFAT3

Protein Interaction Partner: ANKRD28, Dlg4, NR112, MAPK9, MAPK8, JUP, GPR22, CCNG1, LPIN1, CEBPA, GATA4, YWHAZ, MAPK14, CREBBP, YWHAQ,

Protein Size: 902

Molecular Weight: 95 kDa

Gene ID: 4776

NCBI Accession: [NM_004554](#), [NP_004545](#)

UniProt: [Q14934](#)

Pathways: [RTK Signaling](#), [WNT Signaling](#)

Application Details

Application Notes: Optimal working dilutions should be determined experimentally by the investigator.

Comment: Antigen size: 902 AA

Restrictions: For Research Use only

Handling


Format: Liquid

Concentration: Lot specific

Handling


Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images



Western Blotting

Image 1. WB Suggested Anti-NFATC4 Antibody Titration:
0.2-1 ug/ml Positive Control: Transfected 293T



Western Blotting

Image 2. Host: Mouse Target Name: NFATC4 Sample
Tissue: Mouse Pancreas Antibody Dilution: 1ug/ml