

## Datasheet for ABIN2780550

## anti-NR5A2 + LRH1 antibody (N-Term)

# 1 Image



Go to Product page

$\sim$				
	Ive	r\/		٨
$\cup$	' V C	ı vı	$\Box$	٧V

Quantity:	100 μL
Target:	NR5A2 + LRH1 (NR5A2)
Binding Specificity:	N-Term
Reactivity:	Human, Rat, Cow, Guinea Pig, Sheep, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR5A2 + LRH1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human NR5A2
Sequence:	MSSNSDTGDL QESLKHGLTP IVSQFKMVNY SYDEDLEELC PVCGDKVSGY
Predicted Reactivity:	Cow: 100%, Guinea Pig: 86%, Human: 100%, Pig: 92%, Rat: 93%, Sheep: 100%
Characteristics:	This is a rabbit polyclonal antibody against NR5A2. It was validated on Western Blot using a cell
	lysate as a positive control.
Purification:	Protein A purified
Target Details	
Target:	NR5A2 + LRH1 (NR5A2)
Alternative Name:	NR5A2 (NR5A2 Products)


## Target Details

binds to the sequence element 5'-AACGACCGACCTTGAG-3' of the enhancer II of a B virus genes, a critical cis-element of their expression and regulation. It may be table for the liver-specific activity of enhancer II, probably in combination with other syste transcription factors. It is a key regulator of cholesterol 7-alpha-hydroxylase gene expression in liver. It may also contribute to the regulation of pancreas-specific genes of important roles in embryonic development.  Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NR0B2, GPS2, ILF3, COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1,  Size: 541		
hable for the liver-specific activity of enhancer II, probably in combination with other byte transcription factors. It is a key regulator of cholesterol 7-alpha-hydroxylase gene by expression in liver. It may also contribute to the regulation of pancreas-specific genes of important roles in embryonic development.  Important roles in embryonic development.  Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NR0B2, GPS2, ILF3, COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1,  Size: 541		
eyte transcription factors. It is a key regulator of cholesterol 7-alpha-hydroxylase gene expression in liver. It may also contribute to the regulation of pancreas-specific genes of important roles in embryonic development. Important roles in embryonic development. Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NROB2, GPS2, ILF3, COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1, Size: 541		
expression in liver. It may also contribute to the regulation of pancreas-specific genes important roles in embryonic development.  mbols: B1F, B1F2, CPF, FTF, FTZ-F1, FTZ-F1beta, LRH-1, hB1F, hB1F-2, LRH1  Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NR0B2, GPS2, ILF3,  COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D,  MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1,  Size: 541		
mbols: B1F, B1F2, CPF, FTF, FTZ-F1, FTZ-F1beta, LRH-1, hB1F, hB1F-2, LRH1 Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NR0B2, GPS2, ILF3, COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1, Size: 541		
mbols: B1F, B1F2, CPF, FTF, FTZ-F1, FTZ-F1beta, LRH-1, hB1F, hB1F-2, LRH1 Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NR0B2, GPS2, ILF3, COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1, Size: 541		
Interaction Partner: NSD1, NCOA1, CREB1, UBE2I, RAD54L2, NR0B2, GPS2, ILF3, COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1, Size: 541		
COPS5, PNRC1, BCKDK, NRIP1, SRC, NPPA, FTH1, ALB, SUMO1, PPARGC1A, POLR2D, MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1, Size: 541		
MARCD3, PROX1, Ncoa3, Ptpn6, CTNNB1, Size: 541 5860, NP_995582		
Size: 541 5860, NP_995582		
5860, NP_995582		
000482		
Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway		
Optimal working dilutions should be determined experimentally by the investigator.		
Antigen size: 541 AA		
For Research Use only		
Liquid		
Lot specific		
Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 %		
Sodium azide		
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
be handled by trained staff only.		
peated freeze-thaw cycles.		

### Handling

Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Images



#### **Western Blotting**

Image 1. WB Suggested Anti-NR5A2 Antibody Titration:2.5ug/ml Positive Control: HepG2 cell lysate