antibodies -online.com





anti-CIAO1 antibody (C-Term)

2 Images



Go to Product page

Overview

| Quantity: | 100 μL |
|----------------------|---|
| Target: | CIAO1 |
| Binding Specificity: | C-Term |
| Reactivity: | Human, Mouse, Rat, Cow, Dog, Pig, Rabbit, Horse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CIAO1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | The immunogen is a synthetic peptide directed towards the C terminal region of human WDR39 |
|-----------------------|---|
| Sequence: | HSRTIYDIAW CQLTGALATA CGDDAIRVFQ EDPNSDPQQP TFSLTAHLHQ |
| Predicted Reactivity: | Cow: 86%, Dog: 86%, Horse: 86%, Human: 100%, Mouse: 86%, Pig: 86%, Rabbit: 86%, Rat: 86% |
| Characteristics: | This is a rabbit polyclonal antibody against WDR39. It was validated on Western Blot using a cell lysate as a positive control. |
| Purification: | Affinity Purified |

Target Details

| Target: | CIAO1 |
|-------------------|------------------------|
| Alternative Name: | WDR39 (CIAO1 Products) |

Target Details

Handling Advice:


| Background: | WDR39 is a member of the WD40 family of proteins. WDR39 specifically interacts with WT1 |
|---------------------|---|
| | both in vitro and in vivo. This interaction results in a decrease in transcriptional activation |
| | mediated by WT1. WDR39 does not inhibit binding of WT1 to its consensus nucleotide |
| | sequence and does not affect the repression activity of WT1. Thus, WDR39 appears to |
| | specifically modulate the transactivation activity of WT1 and may function to regulate the |
| | physiological functions of WT1 in cell growth and differentiation. |
| | Alias Symbols: CIA1, WDR39 |
| | Protein Interaction Partner: FAM96B, SF1, MEOX1, FAM96A, MMS19, FMNL2, TOE1, DPP9, |
| | FMNL3, CTU1, PPIL4, MAK16, WDR75, NSRP1, EEPD1, WDR61, CDC73, UPF3B, NARFL, |
| | YAE1D1, TYW1, ELP2, ELP3, UCKL1, CDKAL1, DPP8, ELP6, PAF1, PLEKHA5, PHAX, RTEL1, |
| | ELP4, POLA2, ELP5, WDR43, ERP29, GLRX3, CTR9 |
| | Protein Size: 339 |
| Molecular Weight: | 38 kDa |
| Gene ID: | 9391 |
| NCBI Accession: | NM_004804, NP_004795 |
| UniProt: | 076071 |
| Application Details | |
| Application Notes: | Optimal working dilutions should be determined experimentally by the investigator. |
| Comment: | Antigen size: 339 AA |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Format. | Liquid |
| Concentration: | Lot specific |
| Buffer: | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 $\%$ (w/v) sodium azide and 2 $\%$ sucrose. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| | should be handled by trained staff only. |
| | |

Avoid repeated freeze-thaw cycles.

Handling

| Storage: | -20 °C |
|------------------|---|
| Storage Comment: | For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. |

Images



Western Blotting

Image 1. WB Suggested Anti-WDR39 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:312500 Positive Control: Jurkat cell lysate CIAO1 is supported by BioGPS gene expression data to be expressed in Jurkat

Western Blotting

Image 2. WB Suggested Anti-WDR39

Antibody Titration: 0.2-1 µg/mL ELISA Titer: 1:.12500

Positive Control: Jurkat cell lysate

CIAO1 is supported by BioGPS gene expression data to be expressed in Jurkat