



[Go to Product page](#)

Datasheet for ABIN2780940
anti-ZNF432 antibody (C-Term)

1 Image

Overview

Quantity:	100 µL
Target:	ZNF432
Binding Specificity:	C-Term
Reactivity:	Human, Cow, Dog, Horse, Pig, Rabbit, Mouse, Rat, Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF432 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human ZNF432
Sequence:	GKGFTMKSRL IVHQRTHTGE KPFVCSECRK AFSSKRNLIV HQRTHNGNKP
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 91%, Horse: 100%, Human: 100%, Mouse: 91%, Pig: 100%, Rabbit: 100%, Rat: 100%
Characteristics:	This is a rabbit polyclonal antibody against ZNF432. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target:	ZNF432
---------	--------

Target Details

Alternative Name:	ZNF432 (ZNF432 Products)
Background:	ZNF432 belongs to the krueppel C2H2-type zinc-finger protein family. It contains 16 C2H2-type zinc fingers and 1 KRAB domain. ZNF432 may be involved in transcriptional regulation. Alias Symbols: - Protein Size: 652
Molecular Weight:	75 kDa
Gene ID:	9668
NCBI Accession:	NM_014650 , NP_055465
UniProt:	O94892

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 652 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western Blotting

Image 1. WB Suggested Anti-ZNF432 Antibody Titration:
0.2-1 ug/ml Positive Control: Transfected 293T