

# Datasheet for ABIN2780962 anti-HIC2 antibody (C-Term)

## 1 Image



Go to Product page

_				
( )	ve.	rv/	101	Λ

Quantity:	100 μL
Target:	HIC2
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Cow, Horse, Rabbit, Guinea Pig, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIC2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human HIC2
Sequence:	FACDECGMRF TRQYRLTEHM RVHSGEKPYE CQLCGGKFTQ QRNLISHLRM
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit:
	100%, Rat: 100%, Zebrafish: 100%
Characteristics:	This is a rabbit polyclonal antibody against HIC2. It was validated on Western Blot.
Purification:	Affinity Purified
Target Details	
Target:	HIC2
Alternative Name:	HIC2 (HIC2 Products)
, welliative raine.	Tiloz (Tiloz Froducts)

#### Target Details


Background:	HIC2 is a transcriptional repressor.
	Alias Symbols: HRG22, KIAA1020, ZBTB30, ZNF907
	Protein Interaction Partner: HIC2, HIC1, BARD1, LIG4, APP, PDK2, CCNT1,
	Protein Size: 615
Molecular Weight:	66 kDa
Gene ID:	23119
NCBI Accession:	NM_015094, NP_055909
UniProt:	Q96JB3

## Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.	
Comment:	Antigen size: 615 AA	
Restrictions:	For Research Use only	

## Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



#### **Western Blotting**

**Image 1.** WB Suggested Anti-HIC2 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:1562500 Positive Control: THP-1 cell lysate