antibodies .- online.com





anti-AWAT1 antibody (N-Term)



Image



Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | AWAT1 |
| Binding Specificity: | N-Term |
| Reactivity: | Human, Mouse, Rat, Guinea Pig, Cow, Dog, Horse, Rabbit |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AWAT1 antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

Target Details

AWAT1

Target:

| Immunogen: | The immunogen is a synthetic peptide directed towards the N terminal region of human AWAT1 |
|-----------------------|---|
| Sequence: | NWCVWTHIRD YFPITILKTK DLSPEHNYLM GVHPHGLLTF GAFCNFCTEA |
| Predicted Reactivity: | Cow: 92%, Dog: 93%, Guinea Pig: 85%, Horse: 86%, Human: 100%, Mouse: 86%, Rabbit: 86%, Rat: 86% |
| Characteristics: | This is a rabbit polyclonal antibody against DGAT2L3. It was validated on Western Blot using a cell lysate as a positive control. |
| Purification: | Affinity Purified |

Target Details

| Alternative Name: | AWAT1 (AWAT1 Products) |
|-------------------|--|
| Background: | The protein encoded by this gene belongs to the diacylglycerol acyltransferase family. It esterifies long chain (wax) alcohols with acyl-CoA-derived fatty acids to produce wax esters. Wax esters are enriched in sebum, suggesting that this enzyme plays a central role in lipid metabolism in skin. Consistent with this observation, this protein is predominantly expressed in the sebaceous gland of the skin. Alias Symbols: AWAT1, DGA2, DGAT2L3 |
| Molecular Weight: | Protein Size: 328 38 kDa |
| Gene ID: | 158833 |
| NCBI Accession: | NM_001013579, NP_001013597 |
| UniProt: | Q58HT5 |

Application Details

Application Notes:

Precaution of Use:

Handling Advice:

Storage Comment:

Storage:

| Comment: | Antigen size: 328 AA |
|----------------|---|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. |
| Preservative: | Sodium azide |
| | |

should be handled by trained staff only.

Avoid repeated freeze-thaw cycles.


aliquots to prevent freeze-thaw cycles.

-20 °C

Optimal working dilutions should be determined experimentally by the investigator.

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small



Western Blotting

Image 1. WB Suggested Anti-AWAT1 Antibody Titration: 0.2-1 ug/ml Positive Control: Human Stomach