

Datasheet for ABIN2782114
anti-AOC2 antibody (Middle Region)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	AOC2
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Cow, Dog, Horse, Rabbit, Guinea Pig, Rat, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AOC2 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the middle region of human AOC2
Sequence:	QATMVDIHIL VGKGAVQLLP GAVCVFEEAQ GLPLRRHHNY LQNHFYGGGLA
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 83%, Horse: 100%, Human: 100%, Mouse: 100%, Pig: 92%, Rabbit: 100%, Rat: 83%
Characteristics:	This is a rabbit polyclonal antibody against AOC2. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target:	AOC2
---------	------

Target Details

Alternative Name:	AOC2 (AOC2 Products)
Background:	<p>Copper amine oxidases catalyze the oxidative conversion of amines to aldehydes and ammonia in the presence of copper and quinone cofactor. The protein contains several conserved motifs including the active site of amine oxidases and the histidine residues that likely bind copper. It may be a critical modulator of signal transmission in retina, possibly by degrading the biogenic amines dopamine, histamine, and putrescine. Copper amine oxidases catalyze the oxidative conversion of amines to aldehydes and ammonia in the presence of copper and quinone cofactor. This gene shows high sequence similarity to copper amine oxidases from various species ranging from bacteria to mammals. The protein contains several conserved motifs including the active site of amine oxidases and the histidine residues that likely bind copper. It may be a critical modulator of signal transmission in retina, possibly by degrading the biogenic amines dopamine, histamine, and putrescine. This gene may be a candidate gene for hereditary ocular diseases. Alternate splicing results in multiple transcript variants.</p> <p>Alias Symbols: DAO2, RAO</p> <p>Protein Size: 756</p>
Molecular Weight:	84 kDa
Gene ID:	314
NCBI Accession:	NM_009590 , NP_033720
UniProt:	O75106

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 756 AA
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide

Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images



Western Blotting

Image 1. WB Suggested Anti-AOC2 Antibody Titration: 0.2-1 ug/ml Positive Control: Jurkat cell lysate