

Datasheet for ABIN2782179

anti-OMG antibody (Middle Region)






Go to Product page

_				
	IVe	rv	iew	

Quantity:	100 μL	
Target:	OMG	
Binding Specificity:	Middle Region	
Reactivity:	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Rabbit, Goat, Saccharomyces cerevisiae	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This OMG antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	The immunogen is a synthetic peptide directed towards the middle region of human OMG	
Sequence:	NTLRSLEVLN LSSNKLWTVP TNMPSKLHIV DLSNNSLTQI LPGTLINLTN	
Predicted Reactivity:	Cow: 93%, Dog: 100%, Goat: 83%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 79%, Rat: 100%, Yeast: 75%	
Characteristics:	This is a rabbit polyclonal antibody against OMG. It was validated on Western Blot using a cell lysate as a positive control.	
Purification:	Affinity Purified	
Target Details		
Target:	OMG	

Target Details

Alternative Name:	OMG (OMG Products)		
Background:	OMG is a cell adhesion molecule contributing to the interactive process required for myelination		
	in the central nervous system.		
	Alias Symbols: OMGP		
	Protein Interaction Partner: RTN4RL1, RTN4R,		
	Protein Size: 440		
Molecular Weight:	49 kDa		
Gene ID:	4974		
NCBI Accession:	NM_002544, NP_002535		
UniProt:	P23515		
Pathways:	Neurotrophin Signaling Pathway, Regulation of Cell Size		
Application Details			
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.		
Comment:	Antigen size: 440 AA		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	Lot specific		
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
Handling Advice:	Avoid repeated freeze-thaw cycles.		
Storage:	-20 °C		
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small		
	aliquots to prevent freeze-thaw cycles.		



Western Blotting

Image 1. WB Suggested Anti-OMG Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:62500 Positive Control: PANC1 cell lysate