

Datasheet for ABIN2782581
anti-NOC3L antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	NOC3L
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOC3L antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human NOC3L
Sequence:	TLKKYRKEQR KLRQAVKDAV SKKPILENP KEKRP GKRIE REEEEEEEAL
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%
Characteristics:	This is a rabbit polyclonal antibody against NOC3L. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target:	NOC3L
---------	-------

Target Details

Alternative Name:	NOC3L (NOC3L Products)
Background:	<p>NOC3L belongs to the CBF/MAK21 family. It may be required for adipogenesis.</p> <p>Alias Symbols: AD24, C10orf117, FAD24, FLJ12820</p> <p>Protein Interaction Partner: SUMO3, UBC, ZNF622, LIN28A, DICER1, PRKRA, SUMO1, TARBP2, RNF2, SRPK3, EIF2AK2, SIRT7, USP39,</p> <p>Protein Size: 800</p>
Molecular Weight:	92 kDa
Gene ID:	64318
NCBI Accession:	NM_022451 , NP_071896
UniProt:	Q8WTT2

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 800 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western Blotting

Image 1. WB Suggested Anti-NOC3L Antibody Titration:
0.2-1 ug/ml Positive Control: Transfected 293T