



[Go to Product page](#)

Datasheet for ABIN2783094

anti-SNX29 antibody (N-Term)

3 Images

Overview

Quantity:	100 µL
Target:	SNX29 (RUNDC2A)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Horse, Rabbit, Dog, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNX29 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human RUNDC2A
Sequence:	MSGSQNNDKR QLLERLLDA VKQCQIRFGG RKEIASDSDS RVTCLCAQFE
Predicted Reactivity:	Cow: 100%, Dog: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 100%
Characteristics:	This is a rabbit polyclonal antibody against RUNDC2A. It was validated on Western Blot and immunohistochemistry.
Purification:	Protein A purified

Target Details

Target:	SNX29 (RUNDC2A)
---------	-----------------

Target Details

Alternative Name: [RUNDC2A \(RUNDC2A Products\)](#)

Background: The function remains unknown.
Alias Symbols: FLJ12363
Protein Size: 375

Molecular Weight: 42 kDa

Gene ID: 84127

NCBI Accession: [NM_032167](#), [NP_115543](#)

UniProt: [Q9HA26](#)

Application Details

Application Notes: Optimal working dilutions should be determined experimentally by the investigator.

Comment: Antigen size: 375 AA

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: Lot specific

Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling Advice: Avoid repeated freeze-thaw cycles.


Storage: -20 °C

Storage Comment: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.




Immunohistochemistry

Image 1.



Immunohistochemistry

Image 2. Human Intestine



Western Blotting

Image 3. WB Suggested Anti-RUNDC2A Antibody Titration:

5.0ug/ml Positive Control: HepG2 cell lysate