antibodies -online.com





anti-Prohibitin 2 antibody (C-Term)



Images



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	Prohibitin 2 (PHB2)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Rabbit, Guinea Pig, Cow, Dog, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prohibitin 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human PHB2
Sequence:	KLRKIRAAQN ISKTIATSQN RIYLTADNLV LNLQDESFTR GSDSLIKGKK
Predicted Reactivity:	Cow: 100%, Dog: 93%, Guinea Pig: 93%, Human: 100%, Mouse: 100%, Pig: 100%, Rabbit: 100%, Rat: 100%
Characteristics:	This is a rabbit polyclonal antibody against PHB2. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified
Target Details	
Target:	Prohibitin 2 (PHB2)


Target Details

Alternative Name:	PHB2 (PHB2 Products)		
Background:	PHB2 acts as a mediator of transcriptional repression by nuclear hormone receptors via		
	recruitment of histone deacetylases (By similarity). It functions as an estrogen receptor (ER)-		
	selective coregulator that potentiates the inhibitory activities of antiestrogens and represses		
	the activity of estrogens. PHB2 competes with NCOA1 for modulation of ER transcriptional		
	activity. It probably involved in regulating mitochondrial respiration activity and in aging.		
	Alias Symbols: BAP, BCAP37, Bap37, MGC117268, PNAS-141, REA, p22		
	Protein Interaction Partner: HUWE1, IGF1R, HDAC1, ESR1, KIAA1244, NCOR1, SUMO2, STAU1		
	UBC, MDM2, ASB14, ASB17, SUZ12, EED, RNF2, UPF2, ADRB2, CDK2, env, VCAM1, TEAD3,		
	ITGA4, ILF3, FN1, ATF2, C1QBP, AFP, UBL4A, EPAS1, FMNL1, PAXIP1, MDC1, MYOD1, MEF2A		
	FHL2, ECT2, BRCA1, AKT1, Htt, U		
	Protein Size: 299		
Molecular Weight:	33 kDa		
Gene ID:	11331		
NCBI Accession:	NM_007273, NP_009204		
UniProt:	Q99623		
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid		
	Hormone Receptor Signaling		
Application Details			
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.		
Comment:	Antigen size: 299 AA		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	Lot specific		
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 %		
	sucrose.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		

Handling


	should be handled by trained staff only.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-20 °C	
Storage Comment:	rage Comment: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in aliquots to prevent freeze-thaw cycles.	

Images



Western Blotting

Image 1. WB Suggested Anti-PHB2 Antibody Titration: 0.2-1 ug/ml Positive Control: HepG2 cell lysate PHB2 is supported by BioGPS gene expression data to be expressed in HepG2



Western Blotting

Image 2. WB Suggested Anti-PHB2 Antibody

Titration: 1 ug/ml

Positive Control: Rat tissue

See Immunoblot 2 Data and Customer Feedback tab for more information.

250	kDa	—
150	kDa	
100	kDa	
75	kDa	—
50	kDa	
37	kDa	
25	kDa	
		=
15	kDa	
10	kDa	

Western Blotting

Image 3. WB Suggested Anti-PHB2 antibody Titration: 1 ug/mL Sample Type: Human heart

Please check the product details page for more images. Overall 5 images are available for ABIN2783329.