

Datasheet for ABIN2783573
anti-MEK2 antibody (C-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	MEK2 (MAP2K2)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Guinea Pig, Pig, Cow, Horse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MEK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human MAP2K2
Sequence:	IKNPAERADL KMLTNHTFIK RSEVEEVDFAGWLCKTLRLN QPGTPTRTAV
Predicted Reactivity:	Cow: 93%, Dog: 93%, Guinea Pig: 92%, Horse: 93%, Human: 100%, Mouse: 85%, Pig: 93%, Rat: 92%
Characteristics:	This is a rabbit polyclonal antibody against MAP2K2. It was validated on Western Blot and immunohistochemistry.
Purification:	Affinity Purified

Target Details

Target:	MEK2 (MAP2K2)
---------	---------------

Target Details

Alternative Name:	MAP2K2 (MAP2K2 Products)
Background:	<p>MAP2K2 is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in MAP2K2 gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene. The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.</p> <p>Alias Symbols: MAPKK2, MEK2, MKK2, PRKMK2</p> <p>Protein Interaction Partner: CCNDBP1, RAF1, SUMO2, UBC, EGFR, IQGAP1, NFE2L2, YWHAB, MAPK1, PEX14, BRAF, ARAF, FAM65B, ZNF207, LIG4, MAPK8, WDR83, Ksr1, COPS5, MAP2K1, MEPCE, GRIN1, CNKSR1, LAMTOR3, GRIN2D, RGS12, CASP9, MAPK3, MAP2K2, DUSP3, Flna,</p> <p>Protein Size: 400</p>
Molecular Weight:	44 kDa
Gene ID:	5605
NCBI Accession:	NM_030662 , NP_109587
UniProt:	P36507
Pathways:	MAPK Signaling , RTK Signaling , Fc-epsilon Receptor Signaling Pathway , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Toll-Like Receptors Cascades , Signaling of Hepatocyte Growth Factor Receptor , BCR Signaling

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 400 AA
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images




Western Blotting

Image 1. WB Suggested Anti-MAP2K2 Antibody Titration: 0.2-1 ug/ml Positive Control: HepG2 cell lysate There is BioGPS gene expression data showing that MAP2K2 is expressed in HepG2



Immunohistochemistry

Image 2. Human Pancreas



Immunohistochemistry

Image 3. MAP2K2 antibody - C-terminal region Formalin Fixed Paraffin Embedded Tissue: Human Lung Tissue
Observed Staining: Cytoplasm of pneumocytes Primary Antibody Concentration: 1:600 Secondary Antibody: Donkey anti-Rabbit-Cy3 Secondary Antibody Concentration: 1:200
Magnification: 20X Exposure Time: 0.5 - 2.0 sec