

Datasheet for ABIN2783582
anti-TKTL2 antibody (C-Term)[Go to Product page](#)

1 Image

1 Publication

Overview

Quantity:	100 µL
Target:	TKTL2
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Cow, Guinea Pig, Rabbit, Dog, Horse, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TKTL2 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human TKTL2
Sequence:	SSAKATGGRV ITVEDHYREG GIGEA VCAAV SREPDILVHQ LAVSGVPQRG
Predicted Reactivity:	Cow: 93%, Dog: 93%, Guinea Pig: 100%, Horse: 93%, Human: 100%, Mouse: 100%, Rabbit: 93%, Rat: 100%, Zebrafish: 86%
Characteristics:	This is a rabbit polyclonal antibody against TKTL2. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Protein A purified

Target Details

Target:	TKTL2
---------	-------

Target Details

Alternative Name:	TKTL2 (TKTL2 Products)
Background:	The function remains unknown. Alias Symbols: DKFZP434L1717, FLJ32975 Protein Interaction Partner: NFIC, Protein Size: 626
Molecular Weight:	68 kDa
Gene ID:	84076
NCBI Accession:	NM_032136 , NP_115512
UniProt:	Q9H0I9

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 626 AA
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Publications

Product cited in:	Clayton, Daley, MacDonald, Fernandez-Vizarra, Bottegoni, ONeil, Major, Griffin, Zhuang,
-------------------	---

Adewoye, Woolcock, Jones, Goodyear, Elmesmari, Filer, Tennant, Alivernini, Buckley, Pitceathly et al.: "Inflammation causes remodeling of mitochondrial cytochrome c oxidase mediated by the bifunctional gene C15orf48. ..." in: **Science advances**, Vol. 7, Issue 50, pp. eabl5182, (2021) ([PubMed](#)).



Western Blotting

Image 1. WB Suggested Anti-TKTL2 Antibody Titration:
2.5ug/ml Positive Control: Jurkat cell lysate