



[Go to Product page](#)

Datasheet for ABIN2784324
anti-ZNF700 antibody (C-Term)

1 Image

Overview

Quantity:	100 µL
Target:	ZNF700
Binding Specificity:	C-Term
Reactivity:	Human, Cow, Mouse, Rat, Guinea Pig, Horse, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF700 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human ZNF700
Sequence:	CKECEKAFCK FSSFQIHERK HRGEKPYECK HCGNGFTSAK ILQIHARTHI
Predicted Reactivity:	Cow: 86%, Guinea Pig: 86%, Horse: 86%, Human: 100%, Mouse: 86%, Rabbit: 86%, Rat: 92%
Characteristics:	This is a rabbit polyclonal antibody against ZNF700. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target:	ZNF700
---------	--------

Target Details

Alternative Name:	ZNF700 (ZNF700 Products)
Background:	<p>ZNF700 belongs to the krueppel C2H2-type zinc-finger protein family. It contains 18 C2H2-type zinc fingers and 1 KRAB domain. ZNF700 may be involved in transcriptional regulation.</p> <p>Alias Symbols: DKFZp434I1610</p> <p>Protein Interaction Partner: UBC,</p> <p>Protein Size: 742</p>
Molecular Weight:	86 kDa
Gene ID:	90592
NCBI Accession:	NM_144566 , NP_653167
UniProt:	Q9H0M5

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 742 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western Blotting

Image 1. WB Suggested Anti-ZNF700 Antibody Titration:
0.2-1 ug/ml Positive Control: 721_B cell lysate