# antibodies .- online.com







## anti-Pallidin antibody (N-Term)



Image



$\sim$	
( )\/\	rview
$\circ$	

Quantity:	100 μL
Target:	Pallidin (PLDN)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Pig, Dog, Rabbit, Horse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Pallidin antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human PLDN
Sequence:	MSVPGPSSPD GALTRPPYCL EAGEPTPGLS DTSPDEGLIE DLTIEDKAVE
Predicted Reactivity:	Cow: 86%, Dog: 100%, Horse: 92%, Human: 100%, Mouse: 100%, Pig: 100%, Rabbit: 85%, Rat: 93%
Characteristics:	This is a rabbit polyclonal antibody against PLDN. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Pallidin (PLDN) Target:

### **Target Details**


-	
Alternative Name:	PLDN (PLDN Products)
Background:	PLDN may play a role in intracellular vesicle trafficking. It interacts with Syntaxin 13 which
	mediates intracellular membrane fusion.
	Alias Symbols: PA, PALLID, HPS9, PLDN
	Protein Interaction Partner: SYCE3, SKA1, HAUS1, CORO6, CCDC136, CCHCR1, RABGEF1,
	BLOC1S6, NUP62, DCTN2, ABI2, MED21, CDK5R1, STX11, TPM3, SMN2, KRT13, NDN, OLFM2
	IFT20, CCDC22, APP, ELAVL1, PLEKHA5, LURAP1, PLEKHF2, CCDC53, BLOC1S2, BLOC1S4,
	EXOC8, WDYHV1, AGGF1, NDC80, DTNBP1, SNA
	Protein Size: 172
Molecular Weight:	20 kDa
Gene ID:	26258
NCBI Accession:	NM_012388, NP_036520
UniProt:	Q9UL45
Pathways:	Synaptic Vesicle Exocytosis
Application Details	
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 172 AA
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 %
	sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C

#### Handling

Storage Comment:

For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Images**



#### **Western Blotting**

**Image 1.** WB Suggested Anti-PLDN Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:12500 Positive Control: PANC1 cell lysate