

Datasheet for ABIN2787044
anti-PPP2R3C antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	PPP2R3C
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Rabbit, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPP2R3C antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Sequence:	RFYYRLPAED EVLLQKLREE SRAVFLQRKS RELLDNEELQ NLWFLLDKHQ
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 100%
Characteristics:	This is a rabbit polyclonal antibody against Ppp2r3c. It was validated on Western Blot.
Purification:	Affinity Purified

Target Details

Target:	PPP2R3C
Alternative Name:	Ppp2r3c (PPP2R3C Products)
Background:	Ppp2r3c may regulate MCM3AP phosphorylation through phosphatase recruitment. Ppp2r3c

Target Details

	may play a role in the activation-induced cell death of B-cells. Alias Symbols: RGD1309207 Protein Size: 453
Molecular Weight:	49 kDa
Gene ID:	362739
NCBI Accession:	NM_001014196 , NP_001014218
UniProt:	Q6AXZ3
Pathways:	PI3K-Akt Signaling

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 453 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

90 kDa__
65 kDa__
40 kDa__
29 kDa__
22 kDa__



Rabbit Anti-Ppp2r3c Antibody
Catalog Number: ARP57063
Lot Number: QC27962
Lane: Rat Brain Lysate

Antibody Titration: 1.0µg/ml
Gel Concentration: 12%

Western Blotting
Image 1.