

Datasheet for ABIN2787101
anti-CCDC25 antibody (Middle Region)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	CCDC25
Binding Specificity:	Middle Region
Reactivity:	Human, Rat, Mouse, Rabbit, Zebrafish (Danio rerio), Horse, Cow, Dog, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC25 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the middle region of human CCDC25
Sequence:	DLAAEKECRD REERNEKKAQ IQEMKKREKE EMKKKREMDE LRSYSSLMKV
Predicted Reactivity:	Cow: 93%, Dog: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Pig: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 100%
Characteristics:	This is a rabbit polyclonal antibody against CCDC25. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

Target Details

Target:	CCDC25
---------	--------

Target Details

Alternative Name:	CCDC25 (CCDC25 Products)
Background:	The function of this protein remains unknown. Alias Symbols: FLJ10853 Protein Interaction Partner: GMCL1, TRIM23, UBC, HHV8GK18_gp81, ELAVL1, Protein Size: 208
Molecular Weight:	24 kDa
Gene ID:	55246
NCBI Accession:	NM_018246 , NP_060716
UniProt:	Q86WR0

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 208 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western Blotting

Image 1. WB Suggested Anti-CCDC25 Antibody Titration:

0.2-1 ug/ml

Positive Control: Hela cell lysate