

Datasheet for ABIN2787581
anti-PSMC3 antibody (N-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	PSMC3
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Dog, Zebrafish (Danio rerio), Cow, Rabbit, Horse, Guinea Pig, Goat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMC3 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human PSMC3
Sequence:	KDKIKENSEK IKVNKTLPYL VSNVIELLDV DPNDQEEDGA NIDLDSQRKG
Predicted Reactivity:	Cow: 100%, Dog: 100%, Goat: 86%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 93%
Characteristics:	This is a rabbit polyclonal antibody against PSMC3. It was validated on Western Blot.
Purification:	Affinity Purified

Target Details

Target:	PSMC3
Alternative Name:	PSMC3 (PSMC3 Products)

Target Details

Background:	<p>PSMC3 is a subunit of the 26S proteasome. 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex.</p> <p>Alias Symbols: MGC8487, TBP1</p> <p>Protein Interaction Partner: AMOTL2, UBC, PSMD9, PSMC6, PSMC3, KDM1A, HUWE1, SUMO2, PSMD14, MDM2, ASB11, UCHL5, PSMD3, PSMD2, PSMD1, PSMC2, PSMC1, ADRM1, JKAMP, FBXO6, PARK2, BAG3, GADD45A, NOS2, MYC, PSMD13, PSMC4, PSMA8, UBF1, ECD, STIP1, RANBP9, PSMD6, ZFYVE16, USP14, RUVBL1, PSM</p> <p>Protein Size: 439</p>
Molecular Weight:	49 kDa
Gene ID:	5702
NCBI Accession:	NM_002804 , NP_002795
UniProt:	P17980
Pathways:	Mitotic G1-G1/S Phases , DNA Replication , Synthesis of DNA , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 439 AA
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific

Handling

Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images



Western Blotting

Image 1. WB Suggested Anti-PSMC3

Antibody Titration: 0.2-1 µg/mL ELISA Titer: 1:1562500
Positive Control: NCI-H226 cell lysate
PSMC3 is supported by BioGPS gene expression data to be expressed in NCIH226

Western Blotting

Image 2. WB Suggested Anti-PSMC3 Antibody Titration:

0.2-1 ug/ml
ELISA Titer: 1:1562500
Positive Control: NCI-H226 cell lysate