

Datasheet for ABIN2787997

## anti-ANAPC7 antibody (C-Term)



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	100 µL
Target:	ANAPC7
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Guinea Pig, Horse, Rabbit, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ANAPC7 antibody is un-conjugated
Application:	Western Blotting (WB)

#### Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the C-terminal region of Rat Anapc7
Sequence:	NSIREAMVMA NNVYKTLGAN AQLTLTLLATV CLEDPVTQEK AKTLDDKALA
Predicted Reactivity:	Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 93%
Characteristics:	This is a rabbit polyclonal antibody against Anapc7. It was validated on Western Blot.
Purification:	Affinity Purified

#### Target Details

Target:	ANAPC7
Alternative Name:	Anapc7 ( <a href="#">ANAPC7 Products</a> )

## Target Details


Background:	The function of this protein remains unknown. Alias Symbols: - Protein Size: 565
Molecular Weight:	62 kDa
Gene ID:	304490
NCBI Accession:	<a href="#">NM_001107142</a> , <a href="#">NP_001100612</a>
UniProt:	<a href="#">D3ZIT4</a>

## Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 565 AA
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



Western Blotting

**Image 1.** Host: Rabbit  
Target Name: Anapc7  
Sample Tissue: Rat Stomach lysates  
Antibody Dilution: 1.0 µg/mL