

Datasheet for ABIN2788014

anti-UCHL3 antibody (Middle Region)





Go to Product page

_				
()	ve.	rv/	101	Λ

cerevisiae
of Mouse Uchl3
of Mouse Uchl3
of Mouse Uchl3 se: 100%, Rabbit:
e: 100%, Rabbit:
e: 100%, Rabbit:
e: 100%, Rabbit:


Target Details

and promotes insulin-signaling and insulin-induced adipogenesis. It is required for stress-	Background:	Deubiquitinating enzyme (DUB) that controls levels of cellular ubiquitin through processing of ubiquitin precursors and ubiquitinated proteins. Thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8. Uchl3 has a 10-fold preference for Arg and Lys at position P3", and exhibits a preference towards 'Lys-48'-linked Ubiquitin chains. Uchl3 deubiquitinates ENAC in apical compartments, thereby regulating apical membrane recycling. Uchl3 indirectly increases the phosphorylation of IGFIR, AKT and FOXO1	
response retinal, skeletal muscle and germ cell maintenance. Uch13 may be involved in working memory and can hydrolyze UBB(+1), a mutated form of ubiquitin which is not effectively degraded by the proteasome. Alias Symbols: - Protein Interaction Partner: Nedd8, Protein Size: 230 Molecular Weight: 26 kDa Gene ID: 50933 NCBI Accession: NML016723, NP_057932 UniProt: Q9UKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide			
degraded by the proteasome. Alias Symbols: - Protein Interaction Partner: Nedd8, Protein Size: 230 Molecular Weight: 26 kDa Gene ID: 50933 NCBI Accession: NML016723, NPL057932 UniProt: Q9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Preservative: Sodium azide		response retinal, skeletal muscle and germ cell maintenance. Uchl3 may be involved in working	
Alias Symbols: - Protein Interaction Partner. Nedd8, Protein Size: 230 Molecular Weight: 26 kDa Gene ID: 50933 NCBI Accession: NML016723, NPL057932 UniProt: Q9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide		memory and can hydrolyze UBB(+1), a mutated form of ubiquitin which is not effectively	
Protein Interaction Partner: Nedd8, Protein Size: 230 Molecular Weight: 26 kDa Gene ID: 50933 NCBI Accession: NM_016723, NP_057932 UniProt: Q9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide		degraded by the proteasome.	
Molecular Weight: 26 kDa Sene ID: 50933 NCBI Accession: NML016723, NP_057932 UniProt: 09JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide		Alias Symbols: -	
Molecular Weight: 26 kDa Gene ID: 50933 NCBI Accession: NM_016723, NP_057932 UniProt: Q9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide		Protein Interaction Partner: Nedd8,	
Gene ID: 50933 NCBI Accession: NM_016723, NP_057932 UniProt: O9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide		Protein Size: 230	
NCBI Accession: NM_016723, NP_057932 UniProt: Q9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Molecular Weight:	26 kDa	
UniProt: Q9JKB1 Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Gene ID:	50933	
Pathways: Feeding Behaviour, Positive Regulation of fat Cell Differentiation Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	NCBI Accession:	NM_016723, NP_057932	
Application Details Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	UniProt:	Q9JKB1	
Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Pathways:	Feeding Behaviour, Positive Regulation of fat Cell Differentiation	
Comment: Antigen size: 230 AA Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Application Details		
Restrictions: Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.	
Handling Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Comment:	Antigen size: 230 AA	
Format: Liquid Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Restrictions:	For Research Use only	
Concentration: Lot specific Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Handling		
Buffer: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. Preservative: Sodium azide	Format:	Liquid	
sucrose. Preservative: Sodium azide	Concentration:	Lot specific	
Preservative: Sodium azide	Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 %	
		sucrose.	
Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	Preservative:	Sodium azide	
	Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	

Handling

	should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images



Western Blotting

Image 1. Host: Rabbit Target Name: Uchl3 Sample Type: Mouse Testis lysates Antibody Dilution: 1.0ug/ml